



[Go to Product page](#)

Datasheet for ABIN1049501

anti-Glutathione Peroxidase 1 antibody (Internal Region)

1 Image

Overview

Quantity:	100 µg
Target:	Glutathione Peroxidase 1 (GPX1)
Binding Specificity:	Internal Region
Reactivity:	Mouse, Rat
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This Glutathione Peroxidase 1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Purpose:	Gpx1 (mouse, rat)
Immunogen:	Peptide with sequence C-QSTVYAFSARP, from the internal region (near N Terminus) of the protein sequence according to NP_032186.2.
Sequence:	QSTVYAFSAR P
Isotype:	IgG
Cross-Reactivity:	Cow, Mouse, Rat
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Grade:	Verified

Target Details

Target:	Glutathione Peroxidase 1 (GPX1)
Alternative Name:	Gpx1 (GPX1 Products)
Background:	Gpx1, glutathione peroxidase 1, A1195024, AL033363, CGPx, GPx-1, GSHPx-1, Gpx, cellular glutathione peroxidase
Gene ID:	14775, 24404
NCBI Accession:	NP_032186
Pathways:	Thyroid Hormone Synthesis , Sensory Perception of Sound , Skeletal Muscle Fiber Development , Cell RedoxHomeostasis , Negative Regulation of intrinsic apoptotic Signaling , SARS-CoV-2 Protein Interactome


Application Details

Application Notes: Western Blot: Approx 22 kDa band observed in Mouse and Rat Kidney and Liver lysates (calculated MW of 22.3 kDa according to Mouse NP_032186.2 and Rat NP_110453.3).
Recommended concentration: 0.3-1 µg/mL.. Primary incubation 1 hour at room temperature.
Peptide ELISA: antibody detection limit dilution 1:64000.

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.
Storage:	-20 °C
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.



Western Blotting

Image 1. ABIN1049501 (0.3µg/ml) staining of Rat Liver lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.