

Datasheet for ABIN106381  
**anti-Pyranose Oxidase antibody**



[Go to Product page](#)

1 Image

## Overview

Quantity:	100 µg
Target:	Pyranose Oxidase
Reactivity:	E. coli
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This Pyranose Oxidase antibody is un-conjugated
Application:	ELISA

## Product Details

Immunogen:	Pyranose Oxidase [E.coli] Immunogen type: Native
Isotype:	IgG
Characteristics:	Concentration Definition: by UV absorbance at 280 nm

## Target Details

Target:	Pyranose Oxidase
Abstract:	<a href="#">Pyranose Oxidase Products</a>
Background:	Synonyms: P2Ox Pyranose oxidase PROD POD POx EC=1.1.3.10 Pyranose:oxygen 2-oxidoreductase Glucose 2-oxidase FAD-oxidoreductase
UniProt:	<a href="#">Q5G234</a> , <a href="#">P79076</a>

## Application Details

**Application Notes:** PYRANOSE OXIDASE Antibody has been assayed against 1.0 µg of Pyranose Oxidase [E.coli] in a standard sandwich ELISA using Peroxidase conjugated Affinity Purified anti-Goat IgG [H&L] Rabbit (ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:20,000 to 1:100,000 of the reconstitution concentration is suggested for this product.

**Restrictions:** For Research Use only

## Handling

**Format:** Lyophilized

**Reconstitution:** Restore with deionized water (or equivalent)

**Concentration:** 10.0 mg/mL


**Buffer:** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

**Preservative:** Sodium azide

**Precaution of Use:** This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

**Storage:** 4 °C

## Images



### Western Blotting

**Image 1.** Goat anti Pyranose Oxidase antibody was used to detect pyranose oxidase under reducing (R) and non-reducing (NR) conditions. Reduced samples of purified target proteins contained 4% BME and were boiled for 5 minutes. Samples of ~1µg of protein per lane were run by SDS-PAGE. Protein was transferred to nitrocellulose and probed with 1:3000 dilution of primary antibody (ON 4 C in ABIN925618). Detection shown was using Dylight 488 conjugated Donkey anti goat (605-741-125 lot 21094 1:10K in TBS/ABIN925618 1 hr RT) . Images were collected using the BioRad VersaDoc System