

Datasheet for ABIN106701  
**anti-PIPOX antibody (Biotin)**[Go to Product page](#)

## 1 Image

## Overview

Quantity:	100 µg
Target:	PIPOX
Reactivity:	Bacteria
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This PIPOX antibody is conjugated to Biotin
Application:	ELISA, Western Blotting (WB), Immunoprecipitation (IP)

## Product Details

Immunogen:	Sarcosine Oxidase [Microbial] Immunogen type: Native
Isotype:	IgG
Characteristics:	Concentration Definition: by UV absorbance at 280 nm

## Target Details

Target:	PIPOX
Alternative Name:	Sarcosine Oxidase ( <a href="#">PIPOX Products</a> )
Gene ID:	10586229
UniProt:	<a href="#">Q46337</a>

## Application Details


Application Notes:	This product has been assayed against 1.0 µg of Sarcosine Oxidase in a standard capture ELISA using Peroxidase Conjugated Streptavidin and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:4,000 to 1:20,000 of the reconstitution concentration is suggested for this product.  ELISA Dilution: 1:5,000 - 1:20,000 Immunoprecipitation Dilution: 1:100  Western Blot Dilution: 1:500 - 1:5,000
--------------------	---

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

Format:	Lyophilized
Reconstitution:	Restore with deionized water (or equivalent)
Concentration:	10.0 mg/mL
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C

## Images



### Western Blotting

**Image 1.** Goat anti Sarcosine Oxidase antibody was used to detect Sarcosine Oxidase. Samples of ~1 and 0.25 µg of purified Sarcosine Oxidase per lane were run by SDS-PAGE under reducing and non-reducing conditions and reduced samples contained 4% BME and were boiled for 5 minutes. Protein was transferred to nitrocellulose and probed with Goat anti Sarcosine Oxidase (1:5K in MB-0070, ON 4 °C). Primary antibody was detected with Dylight 649 conjugated Donkey anti Goat (605-743-125 1:10K 1.5 hr RT in MB-070 and imaged on the BioRad VersaDoc imaging system.