

# Datasheet for ABIN1078026 anti-FTH1 antibody (AA 1-171)

## 2 Images



#### Go to Product page

_						
	1//	Д	rv	16	٦/	٨
	W	$\vdash$	ΙV	Ιt	٦,	/V

Quantity:	100 μL	
Target:	FTH1	
Binding Specificity:	AA 1-171	
Reactivity:	Sheep	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This FTH1 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

#### **Product Details**

Purpose:	Polyclonal Antibody to Ferritin, Heavy Polypeptide (FTH)	
Immunogen:	FTH (Met1-Leu171)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against FTH. It has been selected for its ability to recognize FTH in immunohistochemical staining and western blotting.	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	


### Target Details

FTH1			
Ferritin, Heavy Polypeptide (FTH1 Products)			
FTH1, FTHL6, PIG15, PLIF, Ferritin Heavy Chain, Apoferritin, Placenta Immunoregulatory Factor, Proliferation-Inducing Protein 15, Cell proliferation-inducing gene 15 protein			
Transition Metal Ion Homeostasis			
Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100 Immunocytochemistry: 5-20 μg/mL,1:25-100 Optimal working dilutions must be determined by end user.			
The thermal stability is described by the loss rate. The loss rate was determined by accelerate thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.			
For Research Use only			
Liquid			
500 μg/mL			
PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.			
Sodium azide			
WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilu azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.			
Avoid repeated freeze-thaw cycles.			
Avoid repeated freeze-thaw cycles.			
Avoid repeated freeze-thaw cycles.  4 °C,-20 °C			
• • • • • • • • • • • • • • • • • • •			

Expiry Date:


12 months

#### **Images**



#### **Western Blotting**

Image 1.



#### **Western Blotting**

Image 2. Western Blot; Sample: Caprine Kidney lysate; Primary Ab: 2μg/ml Rabbit Anti-Ovine FTH Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit lgG Polyclonal Antibody (Catalog: SAA544Rb19)