
# antibodies -online.com







# anti-GSTA1 antibody (AA 2-222)





| Go to | D    |     |      |
|-------|------|-----|------|
|       | Pron | HCL | nane |
|       |      |     |      |

| ( ) | 11/               | IN   | /ie    | A .    |
|-----|-------------------|------|--------|--------|
|     | / // <del> </del> | ۱ ات | /   (− | ' \/\/ |
|     |                   |      |        |        |

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | GSTA1  |
| Binding Specificity: | AA 2-222   |
| Reactivity:          | Rat  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This GSTA1 antibody is un-conjugated   |
| Application:         | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC) |

# **Product Details**

| Immunogen:    | GSTa1 (Ser2-Phe222)  |
|---------------|--|
| Isotype:      | IgG  |
| Specificity:  | The antibody is a rabbit polyclonal antibody raised against GSTa1. It has been selected for its ability to recognize GSTa1 in immunohistochemical staining and western blotting. |
| Purification: | Antigen-specific affinity chromatography   |

# **Target Details**

| Target:  | GSTA1   |
|--|---|
| Alternative Name: Glutathione S Transferase Alpha 1 (GSTa1) (GSTA1 Products) |   |
| Background:  | Alternative Names: GSTA1, GST2, GSTA1-1, GTH1, GST class-alpha member 1, GST HA subunit |

1, GST-epsilon, Glutathione S-transferase A1, N-terminally processed

# **Application Details**

### Application Notes:

Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500
 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

### Comment:


The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37&degC for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions:

For Research Use only


# Handling

| Format:            | Liquid  |
|--------------------|---|
| Concentration:     | Lot specific  |
| Buffer:            | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.                                       |
| Preservative:      | Sodium azide  |
| Precaution of Use: | WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.        |
|                    | Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or |
|                    | eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a         |
|                    | physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute   |
|                    | azide-containing compounds in running water before discarding to avoid accumulation of            |
|                    | potentially explosive deposits in lead or copper plumbing.  |
| Handling Advice:   | Avoid repeated freeze-thaw cycles.  |
| Storage:           | 4 °C  |
| Storage Comment:   | Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.                         |
| Expiry Date:       | 12 months   |




# **Western Blotting**

Image 1.



# Western Blotting

**Image 2.** Western Blot; Sample: Rat Liver lysate; Primary Ab: 2μg/ml Rabbit Anti-Rat GSTa1 Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



# **Immunohistochemistry**

**Image 3.** Used in DAB staining on fromalin fixed paraffinembedded kidney tissue

Please check the product details page for more images. Overall 4 images are available for ABIN1078074.