

Datasheet for ABIN1078449
anti-PAI1 antibody (AA 25-402)



[Go to Product page](#)

3 Images

Overview

Quantity:	100 µL
Target:	PAI1 (SERPINE1)
Binding Specificity:	AA 25-402
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PAI1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Plasminogen Activator Inhibitor 1 (PAI1)
Immunogen:	RPA532Hu01Recombinant Plasminogen Activator Inhibitor 1 (PAI1)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against PAI1. It has been selected for its ability to recognize PAI1 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details


Target:	PAI1 (SERPINE1)
Alternative Name:	PAI1 (SERPINE1 Products)
Background:	SERPINE1, PLANH1, Serpin Peptidase Inhibitor Clade E Member 1, Nexin, Plasminogen Activator Inhibitor Type 1 Serpin E1, Endothelial plasminogen activator inhibitor
Pathways:	p53 Signaling , Cellular Response to Molecule of Bacterial Origin , Carbohydrate Homeostasis , Autophagy , Smooth Muscle Cell Migration

Application Details

Application Notes:	Western blotting: 0.01-2 µg/mL, Immunohistochemistry: 5-20 µg/mL, Immunocytochemistry: 5-20 µg/mL, Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.




Western Blotting

Image 1.



Immunohistochemistry

Image 2. Used in DAB staining on formalin fixed paraffin-embedded Kidney tissue



Immunohistochemistry

Image 3. DAB staining on IHC-P; Samples: Human Breast cancer Tissue