



[Go to Product page](#)

Datasheet for ABIN1078569
anti-TMSB4X antibody (AA 1-44)

3 Images

Overview

Quantity:	100 µL
Target:	TMSB4X
Binding Specificity:	AA 1-44
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

Product Details

Immunogen:	Tb4 (Met1-Ser44)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against Tb4. It has been selected for its ability to recognize Tb4 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

Target Details

Target:	TMSB4X
Alternative Name:	Thymosin Beta 4 (Tb4) (TMSB4X Products)
Background:	Alternative Names: TMSB4X, FX, PTMB4, TB4X, TMSB4, Thymosin,Beta 4,X-Linked, TB-500, Seraspenide, Hematopoietic system regulatory peptide

Target Details

Pathways: [Regulation of Actin Filament Polymerization](#), [Regulation of Muscle Cell Differentiation](#), [Maintenance of Protein Location](#), [Smooth Muscle Cell Migration](#)

Application Details

Application Notes:

- Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: Lot specific

Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

Preservative: Sodium azide


Precaution of Use: **WARNING:** Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: 4 °C


Storage Comment: Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.

Expiry Date: 12 months




Western Blotting

Image 1.



Immunohistochemistry

Image 2. Used in DAB staining on formalin fixed paraffin-embedded kidney tissue



Immunohistochemistry

Image 3. DAB staining on IHC-P; Samples: Rat Pancreas Tissue