

Datasheet for ABIN1078639
anti-UCHL3 antibody (AA 2-230)



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	UCHL3 (Uchl3)
Binding Specificity:	AA 2-230
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This UCHL3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Ubiquitin Carboxyl Terminal Hydrolase L3 (UCHL3)
Immunogen:	UCHL3 (Glu2-Ala230)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against UCHL3. It has been selected for its ability to recognize UCHL3 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Human, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	UCHL3 (Uchl3)
Alternative Name:	UCHL3 (Uchl3 Products)
Background:	Ubiquitin Thiolesterase, Ubiquitin thioesterase L3, Ubiquitin carboxyl-terminal hydrolase isozyme L3
Pathways:	Feeding Behaviour , Positive Regulation of fat Cell Differentiation

Application Details

Application Notes:	Western blotting: 0.2-2 µg/mL, 1:250-2500 Immunohistochemistry: 5-20 µg/mL, 1:25-100 Immunocytochemistry: 5-20 µg/mL, 1:25-100 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	500 µg/mL
Buffer:	PBS, pH 7.4, containing 0.01 % SKL, 1 mM DTT, 5 % Trehalose and Proclin300.
Preservative:	Dithiothreitol (DTT), ProClin
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.

Handling


Expiry Date: 12 months

Images



Western Blotting

Image 1.



Immunohistochemistry

Image 2. Used in DAB staining on formalin fixed paraffin-embedded Heart tissue