.-online.com antibodies

Datasheet for ABIN108081 anti-Urate Oxidase antibody (Biotin)

Image



Overview

Quantity:	100 µg
Target:	Urate Oxidase (UOX)
Reactivity:	Bacillus
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This Urate Oxidase antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunoprecipitation (IP)
Product Details	
Immunogen:	Uricase [Bacillus species]
	Immunogentype:Native
Isotype:	IgG
Characteristics:	Concentration Definition: by UV absorbance at 280 nm
Target Details	
Target:	Urate Oxidase (UOX)
Alternative Name:	Uricase (UOX Products)
Gene ID:	3202700
UniProt:	Q5WBJ3

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN108081 | 09/12/2023 | Copyright antibodies-online. All rights reserved.


Application Details

Application Notes:	This product has been assayed against 1.0 μg of Uricase in a standard capture ELISA using
	Peroxidase Conjugated Streptavidin and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic
	acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:4,000 to
	1:53,000 of the reconstitution concentration is suggested for this product.
	ELISA Dilution: 1:5,000 - 1:20,000 Immunoprecipitation Dilution: 1:100
	Western Blot Dilution: 1:500 - 1:5,000
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Restore with deionized water (or equivalent)
Concentration:	10.0 mg/mL
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C

Images



Western Blotting

Image 1. Goat anti Uricase antibody was used to detect purified Uricase under reducing and non-reducing conditions. Samples of ~1 and 0.25 µg of protein per lane were run by SDS-PAGE and reduced samples of purified Uricase contained 4% BME and were boiled for 5 minutes. Protein was transferred to nitrocellulose and probed with Goat anti Uricase (1:5K in MB-0070, ON 4 C). Primary antibody was detected with Dylight 649 conjugated Donkey anti Goat (605-743-125 1:10K 1.5 hr RT in MB-070 and imaged on the BioRad VersaDoc imaging system.