

Datasheet for ABIN1085836
anti-EIF3G antibody (Subunit G)



[Go to Product page](#)

1 Publication

Overview

Quantity:	100 µL
Target:	EIF3G
Binding Specificity:	Subunit G
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EIF3G antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF)

Product Details

Immunogen:	Human EIF3G
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Antigen Affinity Purified

Target Details

Target:	EIF3G
Alternative Name:	EIF3G (EIF3G Products)
Background:	44 kDa antibody, Anti EIF3S4 antibody, eIF 3 delta antibody, eIF 3 RNA binding subunit antibody, eIF-3 RNA-binding subunit antibody, eIF-3-delta antibody, eIF3 delta antibody, eIF3 p42

Target Details

antibody, eIF3 p44 antibody, eIF3g antibody, EIF3G_HUMAN antibody, EIF3S4 antibody, EIF3S4 antibody antibody, EIF3S4, formerly antibody, Eukaryotic translation initiation factor 3 antibody, Eukaryotic translation initiation factor 3 RNA binding subunit antibody, Eukaryotic translation initiation factor 3 RNA-binding subunit antibody, Eukaryotic translation initiation factor 3 subunit 4 antibody, Eukaryotic translation initiation factor 3 subunit 4 delta 44 kD antibody, Eukaryotic translation initiation factor 3 subunit G antibody, Eukaryotic translation initiation factor 3 subunit p42 antibody, Eukaryotic translation initiation factor 3, subunit 4, formerly antibody, Subunit 4 (delta 44kD) antibody, Subunit 4 delta antibody

UniProt: [Q5RK09](#)

Pathways: [Ribonucleoprotein Complex Subunit Organization](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Comment: Other species are not tested. Please decide the specificity by homology

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: PBS with 0.1 % Sodium Azide, 50 % Glycerol, pH 7.3. -20 °C, Avoid freeze / thaw cycles.

Preservative: Sodium azide

Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling Advice: Avoid freeze and thaw cycles.

Storage: -20 °C, -80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Publications

Product cited in: Sesé, Fuentes, Esteve-Codina, Béjar, McGrail, Thomas, Aasen, Ramón Y Cajal: "Hypoxia-mediated translational activation of ITGB3 in breast cancer cells enhances TGF- β signaling and malignant features in vitro and in vivo." in: **Oncotarget**, Vol. 8, Issue 70, pp. 114856-114876, (2018) ([PubMed](#)).

