# antibodies -online.com





## CLEC4E Protein (AA 41-219) (His tag)



Go to Product page

$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Quantity:	50 μg
Target:	CLEC4E
Protein Characteristics:	AA 41-219
Origin:	Human
Source:	Human Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CLEC4E protein is labelled with His tag.

#### **Product Details**

Purpose:	Recombinant Human CLEC4E/Mincel (C-6His)	
Sequence:	RCVVTFRIFQ TCDEKKFQLP ENFTELSCYN YGSGSVKNCC PLNWEYFQSS CYFFSTDTIS WALSLKNCSA MGAHLVVINS QEEQEFLSYK KPKMREFFIG LSDQVVEGQW QWVDGTPLTK SLSFWDVGEP NNIATLEDCA TMRDSSNPRQ NWNDVTCFLN YFRICEMVGI NPLNKGKSLV DHHHHHH	
Characteristics:	Recombinant Human C-Type Lectin Domain Family 4 Member E/CLEC4E is produced with our mammalian expression system in human cells. The target protein is expressed with sequence (Arg41-Leu219) of Human CLEC4E fused with a polyhistidine tag at the C-terminus.	
Purity:	> 95 % as determined by reducing SDS-PAGE.	
Sterility:	0.2 μm filtered	
Endotoxin Level:	Less than 0.1 ng/μg (1 IEU/μg) as determined by LAL test	

#### Target Details

Target:	CLEC4E	
Alternative Name:	clec4e (CLEC4E Products)	
Sub Type:	Fusionprotein	
Background:	C-Type Lectin Domain Family 4 Member E (CLEC4E) is a 219 amino acid single-pass type II membrane protein that contains one C-type Lectin domain. It is expressed in monocytes, CLEC4E functions as a downstream target of C/EBP beta and is thought to play a role in the inflammatory response, possibly via transcriptional control of C/EBP beta. CLEC4E may play a role in the response to inflammatory stimuli in peritoneal macrophages and may be involved in immune surveillance processes under transcriptional control of CEBPB. Human CLEC4E shares 67% sequence identity with its mouse counterpart, suggesting a similar function between species. CLEC-4E exists as multiple alternatively spliced isoforms that are encoded by a gene which maps to a natural killer gene complex region on human chromosome 12.  Alternative Names: C-Type Lectin Domain Family 4 Member E, C-Type Lectin Superfamily Member 9, Macrophage-Inducible C-Type Lectin, CLEC4E, CLECSF9, MINCLE	
Molecular Weight:	21.7 kDa	
UniProt:	Q9ULY5	

### **Application Details**

For Research Use only	Restrictions:
-----------------------	---------------

#### Handling

Format:	Lyophilized
Reconstitution:	It is not recommended to reconstitute to a concentration less than 100 µg/mL.  Dissolve the lyophilized protein in ddH2O.  Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Buffer:	Lyophilized from a 0.2 µm filtered solution of 20 mM PB, 150 mM NaCl, pH 7.4.
Handling Advice:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting.
Storage:	4 °C/-20 °C/-80 °C
Storage Comment:	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks.  Reconstituted protein solution can be stored at 4-7°C for 2-7 days.  Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Handling
----------

Expiry Date:

3 months