antibodies .- online.com





GNMT Protein (AA 2-294) (His tag)



Go to Product pag

Overview

Quantity:	50 μg
Target:	GNMT
Protein Characteristics:	AA 2-294
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This GNMT protein is labelled with His tag.

Product Details

Purpose:	Recombinant Human Glycine N-Methyltransferase/GNMT (N-6His)	
Sequence:	MGSSHHHHHH SSGLVPRGSH MVDSVYRTRS LGVAAEGLPD QYADGEAARV WQLYIGDTRS RTAEYKAWLL GLLRQHGCQR VLDVACGTGV DSIMLVEEGF SVTSVDASDK MLKYALKERW NRRHEPAFDK WVIEEANWMT LDKDVPQSAE GGFDAVICLG NSFAHLPDCK GDQSEHRLAL KNIASMVRAG GLLVIDHRNY DHILSTGCAP PGKNIYYKSD LTKDVTTSVL IVNNKAHMVT LDYTVQVPGA GQDGSPGLSK FRLSYYPHCL ASFTELLQAA FGGKCQHSVL GDFKPYKPGQ TYIPCYFIHV LKRTD	
Characteristics:	Recombinant Human Glycine N-Methyltransferase/GNMT is produced by our E. coli expression system. The target protein is expressed with sequence (Val2-Asg294) of Human GNMT fused with a 6His tag at the N-terminus.	
Purity:	> 95 % as determined by reducing SDS-PAGE.	
Sterility:	0.2 µm filtered	

Product Details		
Endotoxin Level:	Less than 0.1 ng/μg (1 IEU/μg) as determined by LAL test	
Target Details		
Target:	GNMT	
Alternative Name:	Glycine-N-methyltransferase (GNMT Products)	
Sub Type:	Fusionprotein	
Background:	Glycine N-Methyltransferase (GNMT) is a tetrameric cytosolic protein. GNMT catalyzes the synthesis of N-methylglycine from glycine using S-adenosylmethionine (AdoMet) as the methyl donor. It can affects DNA methylation by regulating the ratio of S-adenosylmethionine to S-adenosylhomocystine, playing an important role in maintaining normal AdoMet levels. GNMT is highly expressed in liver. As a major folate-binding protein, GNMT takes part in the detoxification pathway. Defects in GNMT are the cause of hypermethioninemia. the patients with this deficiency are mild hepatomegaly and chronic elevation of serum transaminases. Alternative Names: Glycine N-Methyltransferase, GNMT	
Molecular Weight:	34.9 kDa	
UniProt:	Q14749	
Pathways:	Cellular Glucan Metabolic Process, Regulation of Carbohydrate Metabolic Process	
Application Details		
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Reconstitution:	It is not recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in ddH20. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.	
Buffer:	Supplied as a 0.2 µm filtered solution of 20 mM TrisHCl, 150 mM NaCl, pH 8.0.	
Handling Advice:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting.	
Storage:	-80 °C	
Storage Comment:	Store at < -20°C, stable for 6 months after receipt. Please minimize freeze-thaw cycles.	

1	1 11	١٠
\vdash	land	unc
	ıarıa	11110

Expiry Date:

6 months