# antibodies -online.com





## TIMP1 Protein (AA 24-207) (His tag)



#### Overview

Quantity:	50 μg
Target:	TIMP1
Protein Characteristics:	AA 24-207
Origin:	Human
Source:	Human Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This TIMP1 protein is labelled with His tag.

### **Product Details**

Purpose:	Recombinant Human Tissue Inhibitor of Metalloproteinases 1/TIMP-1 (C-6His)
Sequence:	CTCVPPHPQT AFCNSDLVIR AKFVGTPEVN QTTLYQRYEI KMTKMYKGFQ ALGDAADIRF VYTPAMESVC GYFHRSHNRS EEFLIAGKLQ DGLLHITTCS FVAPWNSLSL AQRRGFTKTY
	TVGCEECTVF PCLSIPCKLQ SGTHCLWTDQ LLQGSEKGFQ SRHLACLPRE PGLCTWQSLR SQIAVDHHHH HH
Characteristics:	Recombinant Human Tissue Inhibitor of Metalloproteinases 1/TIMP-1 is produced by our mammalian expression system in human cells. The target protein is expressed with sequence (Cys24-Ala207) of HumanTIMP1 fused with a polyhistidine tag at the C-terminus.
Purity:	> 95 % as determined by reducing SDS-PAGE.
Sterility:	0.2 µm filtered
Endotoxin Level:	Less than 0.1 ng/μg (1 IEU/μg) as determined by LAL test

Handling Advice:

Storage Comment:

Storage:

Expiry Date:

Target Details	
Target:	TIMP1
Alternative Name:	Timp-1 (TIMP1 Products)
Sub Type:	Fusionprotein
Background:	Tissue Inhibitor of Metalloproteinases 1 (TIMP-1) complexes with metalloproteinases (such as collagenases) and irreversibly inactivates them by binding to their catalytic zinc cofactor. Also mediates erythropoiesis in vitro, but, unlike IL-3, it is species-specific, stimulating the growth and differentiation of only human and murine erythroid progenitors. Known to act on MMP-1, MMP-2, MMP-3, MMP-7, MMP-8, MMP-9, MMP-10, MMP-11, MMP-12, MMP-13, and MMP-16. TIMP-1 does not act on MMP-14.  Alternative Names: Metalloproteinase Inhibitor 1, Erythroid-Potentiating Activity, EPA, Fibroblast collagenase Inhibitor, Collagenase Inhibitor, Tissue Inhibitor of Metalloproteinases 1, TIMP-1, TIMP1, CLGI, TIMP
Molecular Weight:	21.75 kDa
UniProt:	P01033
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	It is not recommended to reconstitute to a concentration less than 100 µg/mL.  Dissolve the lyophilized protein in ddH2O.  Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Buffer:	Lyophilized from a 0.2 µm filtered solution of 20 mM PB, 150 mM NaCl, pH 7.2.

Reconstituted protein solution can be stored at 4-7°C for 2-7 days.

Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Always centrifuge tubes before opening. Do not mix by vortex or pipetting.

Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks.

4 °C/-20 °C/-80 °C

3 months