

Datasheet for ABIN1101107
anti-alpha Defensin 1 antibody

2 Images

[Go to Product page](#)

Overview

Quantity:	50 µg
Target:	alpha Defensin 1 (DEFA1)
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This alpha Defensin 1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Neutralization (Neut)

Product Details

Brand:	IHC-plus™
Immunogen:	E.coli derived recombinant Human NP-1. Type of Immunogen: Cell extract
Isotype:	IgG
Specificity:	Human NP-1
Purification:	Immunoaffinity purified

Target Details

Target:	alpha Defensin 1 (DEFA1)
Alternative Name:	DEFA1 / HP1 (DEFA1 Products)

Target Details

Background:	Name/Gene ID: DEFA1 Synonyms: DEFA1, DEFA2, Defensin, alpha 2, HP-1, HP1, Myeloid-related sequence, MRS, DEF1, Defensin, alpha 1, HNP-1, Neutrophil defensin 1
Gene ID:	1667
UniProt:	P59665
Pathways:	Cellular Response to Molecule of Bacterial Origin

Application Details

Application Notes:	Approved: ELISA (0.5 - 2 µg/mL), IHC, IHC-P (5 µg/mL), Neut (0.4 - 0.6 µg/mL), WB (0.1 - 0.2 µg/mL) Usage: Immunohistochemistry: LS-B7416 in an IHC-plus antibody, validated by our pathologists for use in IHC against formalin-fixed, paraffin-embedded human tissues (IHC validation procedure). A concentration of 5 µg/mL is recommended. Neutralization: To yield one-half maximal inhibition [ND] of the biological activity of Human NP-1 (100 ng/mL), a concentration of 0.40-0.60 µg/mL of this antibody is required. ELISA: To detect hNP-1 by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.5-2 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Human NP-1 (LS-C104516) as a detection antibody, allows the detection of at least 0.2-0.4 ng/well of recombinant hNP-1. Western Blot: To detect hNP-1 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hNP-1 is 1.5-3 ng/lane, under either reducing or non-reducing conditions.
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Sterile water
Concentration:	Lot specific
Buffer:	Lyophilized from PBS, pH 7.2


Handling

Handling Advice: Avoid freeze/thaw cycles.

Storage: 4 °C, -20 °C


Storage Comment: Store Lyophilized at room temperature up to 1 month
Reconstituted for up to 2 weeks at 2-8°C. Aliquot and freeze at -20°C for long term storage.
Avoid freeze/thaw cycles.

Images



Immunohistochemistry

Image 1. Anti-DEFA1 / NP-1 antibody IHC staining of human spleen. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody ABIN1101107 concentration 5 ug/ml.



Immunohistochemistry

Image 2. Anti-DEFA1 / NP-1 antibody IHC of human spleen. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 5 ug/ml.