

### Datasheet for ABIN1105263

# anti-AGGF1 antibody



Go to Product page

| _ |     |   |    |             |     |
|---|-----|---|----|-------------|-----|
|   | 1// | r | Vİ | $\triangle$ | ۸/  |
|   | V   |   | VI |             | / V |

| Quantity:    | 0.1 mg   |  |
|--------------|--|--|
| Target:      | AGGF1  |  |
| Reactivity:  | Mouse  |  |
| Host:        | Rabbit   |  |
| Clonality:   | Polyclonal   |  |
| Conjugate:   | This AGGF1 antibody is un-conjugated   |  |
| Application: | Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA) |  |

## **Product Details**

| IgG   |
|---|
| The polyclonal antibody reacts with mouse VG5Q, a 84 kDa protein. |
| Species reactivity (tested):Mouse                                 |
| Protein A   |
|   |

## Target Details

| Target:           | AGGF1  |  |
|-------------------|--|--|
| Alternative Name: | AGGF1 / VG5Q (AGGF1 Products)  |  |
| Background:       | VG5Q functions as an angiogenic factor in promoting angiogenesis and suppression of VG5Q   |  |
|                   | expression inhibits vessel formation. Angiogenic factors are critical to the initiation of |  |

angiogenesis and maintenance of the vascular network. Angiogenesis has an essential role in pathological conditions such as cancer and various ischaemic and inflammatory diseases. VG5Q can bind to endothelial cells and promote cell proliferation, suggesting that the protein may act in an autocrine fashion. VG5Q interacts with TWEAK (also known as TNFSF12), another secreted angiogenic factor. VG5Q shows strong expression in blood vessels and is secreted when vessel formation is initiated. VG5Q protein was detected mostly in the cytoplasm and around the nuclei of human microvascular endothelial cells (HMVECs). Furthermore VG5Q is detected in human umbilical vein endothelial cells (HUVECs), human heart fibroblast (HHF) and ovarian cancer cells (OV-3), but low expression was detected in kidney cancer cells (RP-45), HeLa Cells and bladder cancer cells. Synonyms: Angiogenic factor VG5Q, Angiogenic factor with G patch and FHA domains 1, G patch domain-containing protein 7, GPATCH7, Vasculogenesis gene on 5q protein

| Gene ID:        | 66549     |
|-----------------|-----------|
| NCBI Accession: | NP_079906 |
| UniProt:        | Q7TN31    |

#### **Application Details**

| Application Notes: | Optimal working dilution should be determined by the investigator.   |  |  |
|--------------------|--|--|--|
| Restrictions:      | For Research Use only  |  |  |
| Handling           |  |  |  |
| Concentration:     | 0.1 mg/mL  |  |  |
| Buffer:            | PBS, 0.02 % sodium azide, 0.1 % bovine serum albumin   |  |  |
| Preservative:      | Sodium azide   |  |  |
| Precaution of Use: | This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |  |  |
| Storage:           | 4 °C   |  |  |
| Storage Comment:   | Store at 2 - 8 °C.   |  |  |