

Datasheet for ABIN1105356

anti-beta Amyloid antibody (C-Term)



[Go to Product page](#)

1 Image

Overview

Quantity:	0.5 mg
Target:	beta Amyloid (Abeta)
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat, Primate
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This beta Amyloid antibody is un-conjugated
Application:	Western Blotting (WB), Enzyme Immunoassay (EIA)

Product Details

Immunogen:	KLH conjugated to a short peptide with amino acid sequence corresponding to the C- terminal of beta amyloid peptide 42.
Sequence:	MVGGVVIA
Clone:	CA9 3B3
Isotype:	IgG2b
Specificity:	This antibody recognizes the C-terminal sequence of beta amyloid peptide 42 and full length beta amyloid peptide 42. The antibody does not cross react with beta amyloid peptide 40 in dot blotting and ELISA. Cross-reactivity to beta amyloid peptide 43 is about 5 % in ELISA.
Cross-Reactivity (Details):	Species reactivity (tested):Human, Primates, Mouse, Rat.
Purification:	Protein G affinity purified

Target Details

Target:	beta Amyloid (Abeta)
Alternative Name:	Amyloid beta (Abeta Products)
Background:	Amyloid beta peptide 42 (Aβ42) is best known for its role in the formation of senile plaques in the brain of patients with Alzheimer's disease. Aβ42 and Aβ40 are the two major amyloid peptides that are produced after cleavage of amyloid precursor protein by secretases. Aβ42 (42 amino acids) is very fibrillogenic. The beta pleated structure of Aβ42 constitutes the initial and key component of the insoluble amyloid fibril in senile plaque. It is widely accepted that Aβ42 contributes to the pathogenesis of Alzheimer's disease. One proposition is that the deposition of amyloid fibril onto the brain tissue results in Alzheimer's disease. Another is that the neurotoxicity of Aβ42 oligomer is the cause of the disease. Synonyms: Amyloid beta peptide
Gene ID:	351
Pathways:	Inflammasome

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling

Reconstitution:	Restore with Double distilled water to adjust the final concentration to 1.0 mg/mL.
Buffer:	0.01 M PBS pH 7.2
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	-20 °C
Storage Comment:	Store at -20 °C.

Western Blotting

Image 1.

Dot blot- beta Amyloid 1-42 antibody(3B3)

Abeta 40

Abeta 42

