

Datasheet for ABIN1105448

anti-Asialoglycoprotein Receptor 1 antibody (FITC)[Go to Product page](#)

Overview

Quantity:	0.1 mg
Target:	Asialoglycoprotein Receptor 1 (ASGR1)
Reactivity:	Human, Rat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Asialoglycoprotein Receptor 1 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (IF), Enzyme Immunoassay (EIA), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Clone:	8D7
Isotype:	IgG1
Cross-Reactivity (Details):	Species reactivity (tested): Human, Rat.
Purification:	Protein G

Target Details

Target:	Asialoglycoprotein Receptor 1 (ASGR1)
Alternative Name:	ASGR1 (ASGR1 Products)
Background:	The asialoglycoprotein (ASGP) receptor is a transmembrane hepatocellular surface carbohydrate binding glycoproteins lacking terminal sialic acid residues (asialoglycoproteins). Characterization of the ASGP receptor revealed its functional role in the binding, internalization

Target Details

and transport of a wide range of glycoproteins, which have exposed galactose or N-acetylgalactosamine residues, via the process of receptor-mediated endocytosis (RME). The ASGP receptor can bind a variety of important plasma proteins including transport proteins (i.e. transferrin), enzymes such as alkaline phosphatase, immunoglobulins including IgA, apoptotic hepatocytes, fibronectin and platelets. Additionally, the expression of the ASGP receptor has been clinically correlated to the level of hepatic function that is lost during liver diseases related to cancer, viral hepatitis, and cirrhosis. The ASGP receptor consists of major and minor subunits, which in the rat were identified as rat hepatic lectin (RHL) 1 and RHL 2/3, with molecular weights of respectively 42, 49 and 54 kDa. The selective binding (calcium and pH depended) and uptake of terminal galactosyl bearing proteins requires the formation of hetero-oligomers between these major and minor forms. The total ASGP receptor population consisted of two functionally distinct receptor populations, designated State 1 and State 2, which were involved in the endocytosis and intracellular processing of ligands by different pathways. Synonyms: ASGP-R 1, ASGPR 1, Asialoglycoprotein receptor 1, Hepatic lectin H1

UniProt:	P02706
Pathways:	Thyroid Hormone Synthesis

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling

Concentration:	0.1 mg/mL
Buffer:	PBS, 0.02 % Sodium Azide, 1 % BSA
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Store at 2 - 8 °C.