

Datasheet for ABIN1105789 **anti-TNFRSF1A antibody**

[Go to Product page](#)

Overview

Quantity:	0.1 mg
Target:	TNFRSF1A
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TNFRSF1A antibody is un-conjugated
Application:	Western Blotting (WB), Immunoprecipitation (IP), Enzyme Immunoassay (EIA), Functional Studies (Func)

Product Details

Isotype:	IgG
Cross-Reactivity (Details):	Species reactivity (tested):Mouse
Purification:	Protein A

Target Details

Target:	TNFRSF1A
Alternative Name:	CD120a / TNFR1 (TNFRSF1A Products)
Background:	TNF-RI (~55-60 kDa) is present on most cell types and is considered to play a prominent role in cell stimulation by TNF-alpha. TNF-alpha activates inflammatory responses, induces apoptosis, regulates cellular proliferation, and may even promote cancer progression. The effects of TNF-alpha are mediated by TNF-RI and TNF-RII, which have both distinct and overlapping

Target Details

downstream signaling cascades. Induction of cytotoxicity and other functions are mediated largely via TNF-RI. TNF-RI is equally well activated by both the 17 kDa soluble and 26 kDa membrane-bound form, whereas TNF-RII is efficiently activated only by the membrane bound form of TNF-alpha. TNF-RI signaling is initiated when trimeric TNF-alpha binds TNF-RI receptors. Subsequent TNF-RI trimerization promotes the recruitment of a proximal signaling complex composed of TNF Receptor Associated protein with a Death Domain (TRADD), Receptor Interacting Protein (RIP), cellular Inhibitor of Apoptosis Protein 1 (cIAP1), TNF Receptor Associated Factor 2 (TRAF2), and likely TRAF5. Studies with TNF-RI-deficient mice indicate that TNF-RI mediates most of the proliferation, pro-inflammatory, and apoptosis-activating pathways. Synonyms: TNF-R1, TNF-RI, TNFR-I, Tnfrsf1a, Tumor necrosis factor receptor 1, Tumor necrosis factor receptor superfamily member 1A, Tumor necrosis factor receptor type I, p55, p60

Gene ID:	10090
UniProt:	P25118
Pathways:	NF-kappaB Signaling , Apoptosis , Caspase Cascade in Apoptosis , Hepatitis C , Ubiquitin Proteasome Pathway

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling

Concentration:	0.1 mg/mL
Buffer:	PBS, 0.1 % bovine serum albumin
Storage:	4 °C
Storage Comment:	Store at 2 - 8 °C.