

Datasheet for ABIN1105797
anti-TNFRSF1A antibody (FITC)



[Go to Product page](#)

Overview

Quantity:	0.1 mg
Target:	TNFRSF1A
Reactivity:	Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This TNFRSF1A antibody is conjugated to FITC
Application:	Flow Cytometry (FACS), Immunohistochemistry (Frozen Sections) (IHC (fro)), Enzyme Immunoassay (EIA)

Product Details

Clone:	HM104
Isotype:	IgG2a
Cross-Reactivity (Details):	Species reactivity (tested):Mouse.
Purification:	Protein G

Target Details

Target:	TNFRSF1A
Alternative Name:	CD120a / TNFR1 (TNFRSF1A Products)
Background:	TNF-RI (~55-60 kDa) is present on most cell types and is considered to play a prominent role in cell stimulation by TNF-alpha. TNF-alpha activates inflammatory responses, induces apoptosis, regulates cellular proliferation, and may even promote cancer progression. The effects of TNF-

Target Details

alpha are mediated by TNF-RI and TNF-RII, which have both distinct and overlapping downstream signaling cascades. Induction of cytotoxicity and other functions are mediated largely via TNF-RI. TNF-RI is equally well activated by both the 17 kDa soluble and 26 kDa membrane-bound form, whereas TNF-RII is efficiently activated only by the membrane bound form of TNF-alpha. TNF-RI signaling is initiated when trimeric TNF-alpha binds TNF-RI receptors. Subsequent TNF-RI trimerization promotes the recruitment of a proximal signaling complex composed of TNF Receptor Associated protein with a Death Domain (TRADD), Receptor Interacting Protein (RIP), cellular Inhibitor of Apoptosis Protein 1 (cIAP1), TNF Receptor Associated Factor 2 (TRAF2), and likely TRAF5. Studies with TNF-RI-deficient mice indicate that TNF-RI mediates most of the proliferation, pro-inflammatory, and apoptosis-activating pathways. Synonyms: TNF-R1, TNF-RI, TNFR-I, Tnfrsf1a, Tumor necrosis factor receptor 1, Tumor necrosis factor receptor superfamily member 1A, Tumor necrosis factor receptor type I, p55, p60

Gene ID:	10090
UniProt:	P25118
Pathways:	NF-kappaB Signaling , Apoptosis , Caspase Cascade in Apoptosis , Hepatitis C , Ubiquitin Proteasome Pathway

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling

Concentration:	0.1 mg/mL
Buffer:	PBS, 0.02 % Sodium Azide, 1 % BSA
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	-20 °C
Storage Comment:	Prior to reconstitution store at -70 °C. Following reconstitution store the antibody (in aliquots) at

Handling

-20 °C for 6 month.

Expiry Date: 6 months