antibodies -online.com







anti-IL27 antibody





Overview

Quantity:	0.5 mg
Target:	IL27
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This IL27 antibody is un-conjugated
Application:	Flow Cytometry (FACS)

Product Details

Immunogen:	Recombinant Human IL-27.
Clone:	B-G49
Isotype:	lgG2a
Specificity:	Recognizes both natural and recombinant Human IL-27
Cross-Reactivity (Details):	Species reactivity (tested):Human.
Purification:	Ion Exchange Chromatography.
Sterility:	0.22 µm filtered

Target Details

Target:	IL27
Alternative Name:	Interleukin-27 / IL27A (IL27 Products)

Target Details

Preservative:

Storage Comment:

Storage:

rarget Details	
Background:	Like interleukin-23 (IL-23), IL-27 is a recently discovered member of the IL-6/IL-12 family of proinflammatory and immunoregulatory cytokines. It exists as a heterodimer composed of the p40-related protein EBI3 and an IL-12 p35-related protein termed p28. IL-27 is produced after activation by antigen-presenting cells and induces proliferation of naive but not memory CD4+ T-cells. It acts by binding to its receptor WSX-1 and gp130 which results in the activation of a Jak/STAT signaling cascade, suggesting the IL-27 is involved in the regulation of immune
	processes. It has been suggested that IL-27 can also be used as a therapeutic agent against cancer as it can also induce tumor-specific anti-tumor activity mediated through CD8+ T-cells, IFN-gamma, and T-bet.Synonyms: IL-27 subunit alpha, IL-30, IL27-A, IL30, Interleukin-30, p28
Gene ID:	246778
NCBI Accession:	NP_663634
UniProt:	Q8NEV9
Application Details	
Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Concentration:	1.0 mg/mL
Buffer:	Phosphate-buffered saline without Carrier and preservatives. This prodcut is sterile-filtered through 0.22 μ m and treated to remove Endotoxins.

Without preservative

Store the antibody undiluted at 2-8 °C.

4°C

Flow Cytometry

Image 1.



