

Datasheet for ABIN1112051
anti-CD11c antibody (FITC)



[Go to Product page](#)

1 Image **2** Publications

Overview

| | |
|--------------|--|
| Quantity: | 100 tests |
| Target: | CD11c (ITGAX) |
| Reactivity: | Human |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Conjugate: | This CD11c antibody is conjugated to FITC |
| Application: | Flow Cytometry (FACS), Immunofluorescence (IF) |

Product Details

| | |
|------------------|---|
| Clone: | BU-15 |
| Isotype: | IgG1 |
| Characteristics: | Monoclonal Mouse Anti-Human CD11c FITC, is recommended for use in flow cytometry for identification of human receptor for Interleukin-2 (IL-2R) expressing the 55,000 M.W. surface antigen. |

Target Details

| | |
|-------------------|--|
| Target: | CD11c (ITGAX) |
| Alternative Name: | CD11c (ITGAX Products) |
| Background: | This antibody reacts with the X chain of the CD11c/CD18 integrin heterodimer expressed on monocytes and weakly on granulocytes. The antibody stains macrophages in most types of tissues. flow cytometry the antibody labels a T-cell subset and weakly a B-cell subset. |

Target Details

Pathways: [Complement System](#), [Activated T Cell Proliferation](#), [Integrin Complex](#)

Application Details

| | |
|--------------------|--|
| Application Notes: | It is recommended for use in flow cytometry. This reagent is effective for direct immunofluorescence staining of human tissue for flow cytometric analysis using 20 µl/10 ⁶ cells. |
| Comment: | Fluorescein isothiocyanate (Molecular Probes). |
| Sample Collection: | 1. Transfer 100 µl of anticoagulated (EDTA) blood to a 12 x 75 mm polystyrene test tube (10 P6 P cells). 2. Add 20 µl of CD11c FITC and mix gently with a vortex mixer. The 20 µl is a guideline only, the optimal volume should be determined by the individual laboratory. 3. The recommended negative control is a non-reactive FITC-conjugated antibody of the same isotype. 4. Incubate in the dark at room temperature at 4 °C for 30 minutes or at room temperature (20-25 °C) for 15 minutes. 5. Add 1,5 ml of Lysing Solution to each sample and mix gently with a vortex mixer. Incubate for 10 minutes at room temperature in the dark. 6. Centrifuge at 1000 x g for 5 minutes. Gently aspirate the supernatant and discard it leaving approximately 50 µl of fluid. 7. Add 2 ml 0.01 mol/l PBS (It better that it containing 2% bovine serum albumin) and resuspend the cells by using a vortex mixer. 8. Centrifuge at 1000 x g for 5 minutes. Gently aspirate the supernatant and discard it leaving approximately 50 µl of fluid. 9. Resuspend pellet in an appropriate fluid for flow cytometry, e.g. 0.3 ml PBS. The PBS should contain 1% paraformaldehyde (fixative) if samples are not analysed the same day. 10. Analyse on a flow cytometer or store at 2-8 °C in the dark until analysis. Samples can be run up to 24 hours after lysis. |
| Restrictions: | For Research Use only |

Handling

| | |
|--------------------|--|
| Format: | Liquid |
| Buffer: | The conjugate is provided in liquid form in buffer containing 1% bovine serum albumin (BSA) and 0,09% Sodium azide, pH 7.2. |
| Preservative: | Sodium azide |
| Precaution of Use: | 1. The device is not intended for clinical use including diagnosis, prognosis, and monitoring of a disease state, and it must not be used in conjunction with patient records or treatment. 2. This product contains Sodium azide (NaN ₃), a chemical highly toxic in pure form. At product concentrations, though not classified as hazardous, Sodium azide may react with lead and |

Handling

copper plumbing to form highly explosive build-ups of metal azides. Upon disposal, flush with large volumes of water to prevent metal azide build-up in plumbing. 3. As with any product derived from biological sources, proper handling procedures should be used.

Storage: 4 °C

Publications

Product cited in: Marin, Artal, Martin, Carrizo, Andres, Martin-Burriel, Bolea, Sanz, Varona, Godino, Gallego, Garcia-Erce, Villar, Gil, Forner, Cubero, Ros: "Epigenetics modifications and Subclinical Atherosclerosis in Obstructive Sleep Apnea: The EPIOSA study." in: **BMC pulmonary medicine**, Vol. 14, pp. 114, (2015) ([PubMed](#)).

Sánchez, Almeida, Vidriales, López-Berges, García-Marcos, Moro, Corrales, Calmuntia, San Miguel, Orfao: "Incidence of phenotypic aberrations in a series of 467 patients with B chronic lymphoproliferative disorders: basis for the design of specific four-color stainings to be used for minimal residual disease investigation." in: **Leukemia**, Vol. 16, Issue 8, pp. 1460-9, (2002) ([PubMed](#)).

Images




Image 1.