

Datasheet for ABIN1112112
anti-CD31 antibody (APC)



[Go to Product page](#)

1 Image

Overview

Quantity:	100 tests
Target:	CD31 (PECAM1)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD31 antibody is conjugated to APC
Application:	Flow Cytometry (FACS), Immunofluorescence (IF)

Product Details

Clone:	TP1-15
Isotype:	IgG2a
Characteristics:	Monoclonal Mouse Anti-Human CD31 APC is recommended for use in flow cytometry.

Target Details

Target:	CD31 (PECAM1)
Alternative Name:	CD31 (PECAM1 Products)
Background:	Anti-CD31 TP1/15, was included in the Fifth International Workshops on Human Leucocyte Differentiation Antigens. Human CD31 is an adhesion molecule expressed on platelets, endothelial cells, leukocytes and their bone marrow precursors. CD31 plays a role in homophilic adhesion and heterophilic transendothelial migration. Antibody TP1/15 reacts with domain 1 of CD31 and blocks homophilic interaction and heterophilic transendothelial migration. Expressed

Target Details

diffusely by monocytes, neutrophils, and NK cells. Expressed on subsets of T cells, not on circulating B cells.

Pathways: [Regulation of Actin Filament Polymerization](#)

Application Details

Application Notes: It is recommended for use in flow cytometry. This reagent is effective for direct immunofluorescence staining of human tissue for flow cytometric analysis using 20 µl/10⁶ cells.

Comment: Allophycocyanin (Febico, Far East Bio-Tech Co.).

Sample Collection: 1. Transfer 100 µl of anticoagulated (EDTA) blood to a 12 x 75 mm polystyrene test tube (10⁶ cells). 2. Add 20 µl of CD31 APC and mix gently with a vortex mixer. The 20 µl is a guideline only, the optimal volume should be determined by the individual laboratory. 3. The recommended negative control is a non-reactive APC-conjugated antibody of the same isotype. 4. Incubate in the dark at room temperature at 4°C for 30 minutes or at room temperature (20-25 °C) for 15 minutes. 5. Add 100 µl of Lysing Solution to each sample and mix gently with a vortex mixer. Incubate for 10 minutes at room temperature in the dark. 6. Centrifuge at 1000 x g for 5 minutes. Gently aspirate the supernatant and discard it leaving approximately 50 µl of fluid. 7. Add 2 ml 0.01 mol/l PBS (It betters that it containing 2% bovine serum albumin) and resuspend the cells by using a vortex mixer. 8. Centrifuge at 1000 x g for 5 minutes. Gently aspirate the supernatant and discard it leaving approximately 50 µl of fluid. 9. Resuspend pellet in an appropriate fluid for flow cytometry, e.g. 0.3 ml PBS. The PBS should contain 1% paraformaldehyde (fixative) if samples are not analysed the same day. 10. Analyse on a flow cytometer or store at 2-8 °C in the dark until analysis. Samples can be run up to 24 hours after lysis.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: The conjugate is provided in liquid form in buffer containing 1% bovine serum albumin (BSA) and 0,09% Sodium azide, pH 7.2.

Preservative: Sodium azide

Precaution of Use: 1. The device is not intended for clinical use including diagnosis, prognosis, and monitoring of a

disease state, and it must not be used in conjunction with patient records or treatment. 2. This product contains Sodium azide (NaN₃), a chemical highly toxic in pure form. At product concentrations, though not classified as hazardous, Sodium azide may react with lead and copper plumbing to form highly explosive build-ups of metal azides. Upon disposal, flush with large volumes of water to prevent metal azide build-up in plumbing. 3. As with any product derived from biological sources, proper handling procedures should be used.

Storage: 4 °C

Images




Image 1.