## ANTIBODIES ONLINE

# Datasheet for ABIN1112144 anti-CD42b antibody (FITC)

1 Image

1 Publication



Overview

Quantity:	100 tests
Target:	CD42b (GP1BA)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD42b antibody is conjugated to FITC
Application:	Flow Cytometry (FACS), Immunofluorescence (IF)

#### Product Details

Clone:	HIP1
lsotype:	lgG1
Characteristics:	Monoclonal Mouse Anti-Human CD42b FITC, is recommended for use in flow cytometry for
	identification of platelets, thrombocytes and megakaryocytes.

#### Target Details

Target:	CD42b (GP1BA)
Alternative Name:	CD42b (GP1BA Products)
Background:	The antibody is directed against the CD42b antigen, platelet glycoprotein gp1bA, that serves as a receptor for Von Willebrand factor (vWF) and as a high affinity thrombin receptor. AK2 immunblots a band of 130kDa on SDS gels of platelet lysates and gives surface
	immunofluorescence with fixed or unfixed platelets. The antigen is absent in very low levels on

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN1112144 | 11/19/2024 | Copyright antibodies-online. All rights reserved. platelets with the Bernard-Soulier syndrome

### Application Details

Application Notes:	It is recommended for use in flow cytometry. This reagent is effective for direct immunofluorescence staining of human tissue for flow cytometric analysis using 20 µl/10^6 cells.
Comment:	Fluorescein isothiocyanate (Molecular Probes).
Sample Preparation:	1. Transfer 5 µl of anticoagulated (EDTA) blood to a 12 x 75 mm polystyrene test tube (10^6 cells). 2. Add 95 µl of PBS (It better that it containing 2% bovine serum albumin) without Fixative mix gently with a vortex mixer. 3. Add 20 µl of CD42b FITC and mix gently with a vortex mixer. The 20 µl is a guideline only, the optimal volume should be determined by the individual laboratory. 4. The recommended negative control is a non-reactive FITC-conjugated antibody of the same isotype. 5. Incubate in the dark at room temperature at 4°C for 30 minutes or at room temperature (20-25 °C) for 15 minutes. 6. Add 0, 2 ml 0.01 mol/l PBS (It betters that it containing 2% bovine serum albumin) and resuspend the cells by using a vortex mixer. 7. Analyse on a flow cytometer, and the readings of the cytometer are recorded, bearing in the mind that the FSC and SSC light dispersion parameters must be on a logarithmic scale.
Restrictions:	For Research Use only

#### Handling

Format:	Liquid
Buffer:	The conjugate is provided in liquid form in buffer containing 1% bovine serum albumin (BSA) and 0,09% Sodium azide, pH 7.2.
Preservative:	Sodium azide
Precaution of Use:	1. The device is not intended for clinical use including diagnosis, prognosis, and monitoring of a disease state, and it must not be used in conjunction with patient records or treatment. 2. This product contains Sodium azide (NaN3), a chemical highly toxic in pure form. At product concentrations, though not classified as hazardous, Sodium azide may react with lead and copper plumbing to form highly explosive build-ups of metal azides. Upon disposal, flush with large volumes of water to prevent metal azide build-up in plumbing. 3. As with any product derived from biological sources, proper handling procedures should be used.
Storage:	4 °C

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN1112144 | 11/19/2024 | Copyright antibodies-online. All rights reserved. Product cited in:

Williams, Jackson: "Analysis of populations of macrophage-granulocyte progenitor cells
stimulated by activities in mouse lung-conditioned medium." in: Experimental hematology, Vol.
5, Issue 6, pp. 523-34, (1978) (PubMed).

#### Images



