# antibodies -online.com





Datasheet for ABIN1112192

## anti-CD58 antibody (PE)







( )	11	$\sim$	rv		۱ ۸
	1 \ /	┙	I \/	╙	1/1

Quantity:	100 tests
Target:	CD58
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD58 antibody is conjugated to PE
Application:	Flow Cytometry (FACS), Immunofluorescence (IF)
Product Details	

## Product Details

Characteristics:	Monoclonal Mouse Anti-Human CD58 PE is recommended for use in flow cytometry.
Isotype:	lgG1
Clone:	HI58a

## **Target Details**

Target:

CD58

Alternative Name:	CD58 (CD58 Products)
Background:	Recognizes the human CD58 cell surface antigen. CD58 is a cell adhesion molecule, which
	plays a critical role in facilitation of antigen specific recognition through interaction with CD2 on
	T lymphocytes. It is a member of the immunoglobulin superfamily of molecules. CD58 has a
	wide tissue distribution, being present on erythrocytes, platelets, monocytes, a subset of
	lymphocytes, bone marrow cells, epithelium, and endothelial cells. There are approximately

5,000 CD58 molecules on each erythrocyte. There is reduced expression of CD58 on haemopoietic cells in individuals with paroxysmal nocturnal haemoglobinuria.

### **Application Details**

#### Application Notes:

It is recommended for use in flow cytometry. This reagent is effective for direct immunofluorescence staining of human tissue for flow cytometric analysis using 20  $\mu$ I/10<sup>6</sup> cells.

#### Comment:

R-Phycoerythrin (Europa Bioproducts, Ely, Cambridge).

#### Sample Collection:

1. Transfer 100  $\mu$ l of anticoagulated (EDTA) blood to a 12 x 75 mm polystyrene test tube (10^6 cells). 2. Add 20  $\mu$ l of CD58 PE and mix gently with a vortex mixer. The 20  $\mu$ l is a guideline only, the optimal volume should be determined by the individual laboratory. 3. The recommended negative control is a non-reactive PE-conjugated antibody of the same isotype. 4. Incubate in the dark at room temperature at 4°C for 30 minutes or at room temperature (20-25 °C) for 15 minutes. 5. Add 1,5 ml of Lysing Solution to each sample and mix gently with a vortex mixer. Incubate for 10 minutes at room temperature in the dark. 6. Centrifuge at 1000 x g for 5 minutes. Gently aspirate the supernatant and discard it leaving approximately 50  $\mu$ l of fluid. 7. Add 2 ml 0.01 mol/l PBS (It betters that it containing 2% bovine serum albumin) and resuspend the cells by using a vortex mixer. 8. Centrifuge at 1000 x g for 5 minutes. Gently aspirate the supernatant and discard it leaving approximately 50  $\mu$ l of fluid. 9. Resuspend pellet in an appropriate fluid for flow cytometry, e.g. 0.3 ml PBS. The PBS should contain 1% paraformaldehyde (fixative) if samples are not analysed the same day. 10. Analyse on a flow cytometer or store at 2-8 °C in the dark until analysis. Samples can be run up to 24 hours after lysis.

#### Restrictions:

For Research Use only

## Handling


Format:	Liquid
Buffer:	The conjugate is provided in liquid form in buffer containing 1% bovine serum albumin (BSA) and 0,09% Sodium azide, pH 7.2.
Preservative:	Sodium azide
Precaution of Use:	1. The device is not intended for clinical use including diagnosis, prognosis, and monitoring of a disease state, and it must not be used in conjunction with patient records or treatment. 2. This product contains Sodium azide (NaN3), a chemical highly toxic in pure form. At product

concentrations, though not classified as hazardous, Sodium azide may react with lead and copper plumbing to form highly explosive build-ups of metal azides. Upon disposal, flush with large volumes of water to prevent metal azide build-up in plumbing. 3. As with any product derived from biological sources, proper handling procedures should be used.

Storage:

4°C

## Validation report #028889 for Immunofluorescence (IF)



#### Image 1.