

Datasheet for ABIN116367
anti-IL-3 beta antibody (Biotin)



[Go to Product page](#)

Overview

Quantity:	25 µg
Target:	IL-3 beta
Reactivity:	Rat
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This IL-3 beta antibody is conjugated to Biotin
Application:	Western Blotting (WB), Enzyme Immunoassay (EIA)

Product Details

Immunogen:	Highly pure (>98%) recombinant rIL-3 beta.
Specificity:	Rat Interleukin-3 beta
Purification:	Affinity chromatography.

Target Details

Target:	IL-3 beta
Alternative Name:	Interleukin-3 beta / IL3 beta (IL-3 beta Products)
Background:	<p>IL3 acts in hematopoiesis by controlling the production, differentiation, and function of two related white cell populations of the blood, the granulocytes and the monocytes/macrophages. It induces granulocytes, macrophages, mast cells, stem cells, erythroid cells, eosinophils and megakaryocytes and is expressed by activated T cells, mast cells, and natural killer cells. Synonyms: IL-3 beta, IL3B</p>

Target Details

Gene ID:	24495
NCBI Accession:	NP_113701
UniProt:	P97688

Application Details

Application Notes:	<p>ELISA: To detect rat IL-3 beta by direct ELISA (using 100 l/well antibody solution) this antibody can be used at a concentration of 0.15 - 0.30 µg/mL. Used in conjunction with compatible secondary reagents, allows the detection of at least 0.2 ng/well of recombinant rat IL-3 beta.</p> <p>Western Blot: To detect rat IL-3 beta by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant rat IL-3 beta is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.</p> <p>Other applications not tested.</p> <p>Optimal dilutions are dependent on conditions and should be determined by the user.</p>
Restrictions:	For Research Use only

Handling

Reconstitution:	Restore in sterile PBS containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL.
Buffer:	PBS, pH 7.2 without preservatives.
Preservative:	Without preservative
Handling Advice:	Avoid repeated freezing and thawing. Centrifuge vial prior to opening!
Storage:	4 °C/-20 °C
Storage Comment:	Store the antibody prior to reconstitution at -20 °C. Following reconstitution the antibody can be stored at 2-8 °C for one month or at -20 °C for longer.