antibodies .- online.com





anti-SIRT2 antibody



Image



Go to Product page

()	1 /	\sim	rv	11/	11	Α
	1//	┙	I \/	16	٦,	/\

Quantity:	100 μg
Target:	SIRT2
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), ELISA

Product Details

Immunogen:	Recombinant human sirtuin 2.
Specificity:	Recognizes human sirtuin 2. Detects a band of ~45 kDa by Western blot.
Cross-Reactivity:	Human
Sterility:	0.2 µm filtered

Target Details

Target:	SIRT2
Alternative Name:	Sirtuin 2 (SIRT2 Products)
Background:	Sirtuin-2 is a NAD-dependent protein deacetylase, which deacetylates the 'Lys-40' of a-tubulin. It is involved in the control of mitotic exit in the cell cycle. Furthermore, a variety of age-associated diseases such as type 2 diabetes, obesity, osteoporosis, and Alzheimer's disease
	are associated with sirtuins. Selective inhibition of Sirtuin-2 achieves neuroprotection and identifies Sirtuin-2 to be a potential therapeutic avenue in Huntington's and Parkinson's disease.

Target Details

UniProt: Q8IXJ6


Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	0.2µm-filtered solution in PBS, pH 7.4. Contains no preservatives.
Preservative:	Without preservative
Storage:	4 °C,-20 °C
Storage Comment:	Short Term Storage: +4°C
	Long Term Storage: -20°C
	Stable for at least 6 months after receipt when stored at -20°C.
Expiry Date:	6 months

Images



Western Blotting

Image 1. Western blot analysis using anti-Sirtuin 2 (human), pAb at 1:4'000 dilution. 1: Human Sirtuin 2 (His-tagged). 2: Human FGF21 (His-tagged) (negative control).