antibodies -online.com





Go to Product page

Datasheet for ABIN1171845

anti-SLC13A2 antibody

Overview

| Quantity: | 100 μL |
|--------------|---|
| Target: | SLC13A2 |
| Reactivity: | Pig |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This SLC13A2 antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Immunohistochemistry (Paraffinembedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |

Product Details

| Immunogen: | The antibody is a rabbit polyclonal antibody raised against MUC. |
|---------------|---|
| Isotype: | IgG |
| Specificity: | It has been selected for its ability to recognize MUC in immunohistochemical staining and Western blotting. |
| Purification: | Affinity Chromatography |

Target Details

| Target: | SLC13A2 |
|-------------------|--------------------------------|
| Alternative Name: | Mucin (MUC) (SLC13A2 Products) |

Application Details

Expiry Date:

12 months

| Restrictions: | For Research Use only |
|--------------------|---|
| | |
| Handling | |
| Format: | Lyophilized |
| Concentration: | Lot specific |
| Buffer: | Supplied as solution form in PBS, pH7.4, containing 0.02 % NaN3, 50 % glycerol. |
| Preservative: | Sodium azide |
| Precaution of Use: | WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. |
| | Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or |
| | eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a |
| | physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute |
| | azide-containing compounds in running water before discarding to avoid accumulation of |
| | potentially explosive deposits in lead or copper plumbing. |
| Handling Advice: | Avoid repeated freeze/thaw cycles |
| Storage: | 4 °C |
| Storage Comment: | Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months. |
| | |