

Datasheet for ABIN1172008
anti-PROCR antibody (Biotin)



[Go to Product page](#)

Overview

Quantity:	200 µL
Target:	PROCR
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PROCR antibody is conjugated to Biotin
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

Product Details

Purpose:	Biotin-Linked Polyclonal Antibody to Endothelial protein C receptor (EPCR)
Immunogen:	The antibody is a rabbit polyclonal antibody raised against PROCR conjugated to biotin.
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against EPCR. It has been selected for its ability to recognize EPCR in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	PROCR
Alternative Name:	Endothelial protein C receptor (PROCR Products)
Background:	CD201, PROCR, CCCA ,CCD41, BA4204.2, EPCR, Endothelial Protein C Receptor, Cell Cycle,

Target Details

Centrosome-Associated Protein, Activated Protein C Receptor

Application Details

Application Notes:	Western blotting: 0.2-2 µg/mL, 1:250-2500 Immunohistochemistry: 5-20 µg/mL, 1:25-100 Immunocytochemistry: 5-20 µg/mL, 1:25-100 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	500 µg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze/thaw cycles
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	12 months