

Datasheet for ABIN1172095
anti-IGFBP3 antibody



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	IGFBP3
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IGFBP3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Insulin Like Growth Factor Binding Protein 3 (IGFBP3)
Immunogen:	RPA054Hu01Recombinant Insulin Like Growth Factor Binding Protein 3 (IGFBP3)
Sequence:	MGHHHHHHSG S-GNAESEED RSAGSVESPS VSSTHRVSDP KFHPLHSKII IIKKGHAKDS QRYKVDYESQ STDTQNFSS SKRETEYGPC RREMEDTLNH LKFLNVLSPR GVHIPNCDKK GFYKKKQCRP SKGRKRGFCW CVDKYGQPLP GYTT
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against IGFBP3. It has been selected for its ability to recognize IGFBP3 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	IGFBP3
Alternative Name:	IGFBP3 (IGFBP3 Products)
Background:	BP53, IBP3, Growth Hormone-Dependent Binding Protein, Acid Stable Subunit Of The 140 K IGF Complex
Pathways:	Myometrial Relaxation and Contraction , Regulation of Muscle Cell Differentiation , Skeletal Muscle Fiber Development , Regulation of Carbohydrate Metabolic Process , Autophagy , Smooth Muscle Cell Migration , Growth Factor Binding

Application Details

Application Notes:	Western blotting: 0.01-3 µg/mL, Immunohistochemistry: 5-20 µg/mL, Immunocytochemistry: 5-20 µg/mL, Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling


Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze/thaw cycles
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without

Handling

detectable loss of activity. Avoid repeated freeze-thaw cycles.


Expiry Date: 12 months

Images



Western Blotting

Image 1.



Immunohistochemistry

Image 2. Figure.DAB staining on IHC-P. Samples: Human Tissue