

Datasheet for ABIN1172151 anti-MGEA5 antibody (AA 19-222)

2 Images




Go to Product page

()	ve	rvi	6	W
\sim	v C	1 V I	\sim	v v

Overview		
Quantity:	100 μL	
Target:	MGEA5	
Binding Specificity:	AA 19-222	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This MGEA5 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)	
Product Details		
Immunogen:	NAGase (Ala19-Thr222)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against NAGase. It has been selected for its ability to recognize NAGase in immunohistochemical staining and western blotting.	
Purification:	Antigen-specific affinity chromatography	
Target Details		
Target:	MGEA5	
	N-Acetyl beta-D-Glucosaminidase (NAGase) (MGEA5 Products)	
Alternative Name:	N-Acetyl beta-D-Glucosaminidase (NAGase) (MGEA5 Products)	
Alternative Name: Background:	N-Acetyl beta-D-Glucosaminidase (NAGase) (MGEA5 Products) Alternative Names: MGEA5, MEA5, NCOAT, Hexosaminidase C, N-Acetylhexosaminidase,	


Target Details

	Hyaluronidase, Meningioma-Expressed Antigen 5, Nuclear Cytoplasmic O-GlcNAcase And Acetyltransferase	
Pathways:	Positive Regulation of Peptide Hormone Secretion, Regulation of Carbohydrate Metabolic Process	
Application Details		
Application Notes:	Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	Lot specific	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	4 °C	
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.	
Expiry Date:	12 months	



Western Blotting

Image 1.



Immunohistochemistry

Image 2. Figure.DAB staining on IHC-P. Samples: Rat Tissue