

Datasheet for ABIN1173425 anti-Neuropilin 1 antibody (Biotin)



200 μL
Neuropilin 1 (NRP1)
Mouse
Rabbit
Polyclonal
This Neuropilin 1 antibody is conjugated to Biotin
Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Biotin-Linked Polyclonal Antibody to Neuropilin 1 (NRP1)
The antibody is a rabbit polyclonal antibody raised against NRP1 conjugated to biotin.
IgG
The antibody is a rabbit polyclonal antibody raised against NRP1. It has been selected for its ability to recognize NRP1 in immunohistochemical staining and western blotting.
Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Neuropilin 1 (NRP1)
Neuropilin 1 (NRP1 Products)
CD304, BDCA4, VEGF165R, Vascular endothelial cell growth factor 165 receptor

Target Details

Pathways:	Regulation of Cell Size, Signaling Events mediated by VEGFR1 and VEGFR2, Smooth Muscle Cell Migration, Platelet-derived growth Factor Receptor Signaling, VEGFR1 Specific Signals
Application Details	
Application Notes:	Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100 Immunocytochemistry: 5-20 μg/mL,1:25-100 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	500 μg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze/thaw cycles
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	12 months