

### Datasheet for ABIN1173541

## anti-Glucocorticoid Receptor antibody (AA 622-755)

2 Images



Go to Product page

_				
()	ve.	rv/	101	Λ

Quantity:	100 μL	
Target:	Glucocorticoid Receptor (NR3C1)	
Binding Specificity:	AA 622-755	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Glucocorticoid Receptor antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP),	
	Immunocytochemistry (ICC)	
Product Details		
Product Details  Purpose:	Polyclonal Antibody to Glucocorticoid Receptor Alpha (GRa)	
	Polyclonal Antibody to Glucocorticoid Receptor Alpha (GRa)  RPA774Mu01Recombinant Glucocorticoid Receptor Alpha (GRa)	
Purpose:		
Purpose: Immunogen:	RPA774Mu01Recombinant Glucocorticoid Receptor Alpha (GRa)	
Purpose: Immunogen: Isotype:	RPA774Mu01Recombinant Glucocorticoid Receptor Alpha (GRa)	
Purpose: Immunogen: Isotype:	RPA774Mu01Recombinant Glucocorticoid Receptor Alpha (GRa)  IgG  The antibody is a rabbit polyclonal antibody raised against GRa. It has been selected for its	
Purpose: Immunogen: Isotype: Specificity:	RPA774Mu01Recombinant Glucocorticoid Receptor Alpha (GRa)  IgG  The antibody is a rabbit polyclonal antibody raised against GRa. It has been selected for its ability to recognize GRa in immunohistochemical staining and western blotting.	

### **Target Details**

l arget Details		
Alternative Name:		
Background:		
Pathways:	Nuclear Receptor Transcription Pathway, Intracellular Steroid Hormone Receptor Signaling	
	Pathway, Steroid Hormone Mediated Signaling Pathway, Regulation of Intracellular Steroid	
	Hormone Receptor Signaling, Regulation of Hormone Metabolic Process, Regulation of	
	Hormone Biosynthetic Process, Regulation of Muscle Cell Differentiation, Regulation of	
	Carbohydrate Metabolic Process	
Application Details		
Application Notes:	Western blotting: 0.01-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-	
	20 μg/mL,Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
	date under appropriate storage condition.	

For Research Use only

# Restrictions:


Handling		
Format:	Liquid	
Concentration:	0.5 mg/mL	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	

detectable loss of activity. Avoid repeated freeze-thaw cycles.

Expiry Date:


12 months

### **Images**



### **Western Blotting**

Image 1.



### **Immunohistochemistry**

**Image 2.** Figure.DAB staining on IHC-P. Samples: Mouse Tissue