# antibodies -online.com







# anti-PYGL antibody (AA 341-509)





$() \cup \cap \cap \cup \cap \cup$	
Overview	

Quantity:	100 μL
Target:	PYGL
Binding Specificity:	AA 341-509
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

## **Product Details**


Immunogen:	PYGL (Thr341-Gly509)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against PYGL. It has been selected for its ability to recognize PYGL in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

## **Target Details**

Target:	PYGL
Alternative Name:	Glycogen Phosphorylase, Liver (PYGL) (PYGL Products)
Background:	Alternative Names: GPLL, GPBB, Hers Disease, Glycogen Storage Disease Type VI
Pathways:	Carbohydrate Homeostasis, Cellular Glucan Metabolic Process

# **Application Details**

Application Notes:	<ul> <li>Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500         Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.     </li> </ul>
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37&degC for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	4 °C
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.
Expiry Date:	12 months



### **Western Blotting**

#### Image 1.

## **Western Blotting**

Image 2. Knockout Varification: ;Lane 1: Wild-type Hepg2 cell lysate; ;Lane 2: PYGL knockout Hepg2 cell lysate; ;Predicted MW: 97kDa ;Observed MW: 100kDa;Primary Ab: 1μg/ml Rabbit Anti-Rat PYGL Ab;Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody;(Catalog: SAA544Rb19)

### **Western Blotting**

**Image 3.** Western Blot; Sample: Rat Liver lysate; Primary Ab:  $1\mu g/ml$  Rabbit Anti-Rat PYGL Antibody Second Ab:  $0.2\mu g/mL$  HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)

Please check the product details page for more images. Overall 4 images are available for ABIN1173786.