

Datasheet for ABIN1174356

anti-Fibrinogen beta Chain antibody (AA 33-479)[Go to Product page](#)

4 Images

Overview

Quantity:	100 µL
Target:	Fibrinogen beta Chain (FGB)
Binding Specificity:	AA 33-479
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Fibrinogen beta Chain antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

Product Details

Immunogen:	FGb (Gly33-Gln479)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against FGb. It has been selected for its ability to recognize FGb in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

Target Details


Target:	Fibrinogen beta Chain (FGB)
Alternative Name:	Fibrinogen Beta (FGb) (FGB Products)
Background:	Alternative Names: FG-B, Fibrinopeptide B

Application Details

Application Notes:	<ul style="list-style-type: none">Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	4 °C
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.
Expiry Date:	12 months




Western Blotting

Image 1.



Western Blotting

Image 2. Western Blot; Sample: Rat Spleen lysate; Primary Ab: 1µg/ml Rabbit Anti-Rat FGb Antibody Second Ab: 0.2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



Immunohistochemistry

Image 3. Used in DAB staining on formalin fixed paraffin-embedded Liver tissue

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN1174356.