

Datasheet for ABIN1174692 anti-CSF1R antibody (AA 768-928)

2 Images



Go to Product page

Overview

Quantity:	100 μL
Target:	CSF1R
Binding Specificity:	AA 768-928
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CSF1R antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP)

Product Details

Purpose:	Polyclonal Antibody to Colony Stimulating Factor Receptor, Macrophage (MCSFR)
Immunogen:	RPB484Hu01Recombinant Colony Stimulating Factor Receptor, Macrophage (MCSFR)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against MCSFR. It has been selected for its ability to recognize MCSFR in immunohistochemical staining and western blotting.
Cross-Reactivity:	Pig
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography


Target Details

Target:	CSF1R
Alternative Name:	MCSFR (CSF1R Products)
Background:	CD115, MCFSR, M-CSFR, CSF1R, CSF1-R, C-FMS, CSFR, FIM2, FMS, Colony Stimulating Factor 1 Receptor, Formerly McDonough Feline Sarcoma Viral(v-Fms)oncogene Homolog
Pathways:	RTK Signaling, Inositol Metabolic Process, Cell-Cell Junction Organization
Application Details	
Application Notes:	Western blotting: 0.01-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-20 μg/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.31 mg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.

Expiry Date:


12 months

Images



Western Blotting

Image 1.



Immunohistochemistry

Image 2. Used in DAB staining on fromalin fixed paraffinembedded Liver tissue