

Datasheet for ABIN1174698
anti-CSF1R antibody (AA 734-900)



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	CSF1R
Binding Specificity:	AA 734-900
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CSF1R antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP)

Product Details

Purpose:	Polyclonal Antibody to Colony Stimulating Factor Receptor, Macrophage (MCSFR)
Immunogen:	RPB484Ra01Recombinant Colony Stimulating Factor Receptor, Macrophage (MCSFR)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against MCSFR. It has been selected for its ability to recognize MCSFR in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	CSF1R
Alternative Name:	MCSFR (CSF1R Products)
Background:	CD115, MCFSR, M-CSFR, CSF1R, CSF1-R, C-FMS, CSFR, FIM2, FMS, Colony Stimulating Factor 1 Receptor, Formerly McDonough Feline Sarcoma Viral(v-Fms)oncogene Homolog
Pathways:	RTK Signaling , Inositol Metabolic Process , Cell-Cell Junction Organization

Application Details

Application Notes:	Western blotting: 0.5-2 µg/mL, Immunohistochemistry: 5-20 µg/mL, Immunocytochemistry: 5-20 µg/mL, Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	500 µg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.

Handling


Expiry Date: 12 months

Images



Western Blotting

Image 1.



Immunohistochemistry

Image 2. DAB staining on IHC-P; Samples: Rat Brain Tissue