

Datasheet for ABIN1174718

anti-TNFRSF1A antibody (AA 43-194)





Go to Product page

()	ve	V /	-	1 A
	\cup	1 \/	-	1/1
\sim	' V C	1 V	ı	v v

Quantity:	100 μL
Target:	TNFRSF1A
Binding Specificity:	AA 43-194
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TNFRSF1A antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Tumor Necrosis Factor Receptor 1 (TNFR1)
Immunogen:	RPB499Ra01Recombinant Tumor Necrosis Factor Receptor 1 (TNFR1)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against TNFR1. It has been selected for its ability to recognize TNFR1 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details


Target:	TNFRSF1A
Alternative Name:	TNFR1 (TNFRSF1A Products)
Background:	CD120A, P55, TNFRSF1A, TBP1, FPF, TNF-R, TNF-R-I, TNF-R55, TNFAR, TNFR55, TNFR60, P55-R, P60, Tumor necrosis factor receptor 1, Tumor necrosis factor-binding protein 1
Pathways:	NF-kappaB Signaling, Apoptosis, Caspase Cascade in Apoptosis, Hepatitis C, Ubiquitin Proteasome Pathway
Application Details	
Application Notes:	Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100 Immunocytochemistry: 5-20 μg/mL,1:25-100 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Restrictions: Handling	For Research Use only
	For Research Use only Liquid
Handling	
Handling Format:	Liquid
Handling Format: Concentration:	Liquid 0.29 mg/mL
Handling Format: Concentration: Buffer:	Liquid 0.29 mg/mL PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Handling Format: Concentration: Buffer: Preservative:	Liquid 0.29 mg/mL PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. ProClin, Sodium azide WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of
Handling Format: Concentration: Buffer: Preservative: Precaution of Use:	Liquid 0.29 mg/mL PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. ProClin, Sodium azide WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.

detectable loss of activity. Avoid repeated freeze-thaw cycles.

Expiry Date:

12 months

Images




Western Blotting

Image 1.



Western Blotting

Image 2. Western Blot; Sample: Rat Cerebrum lysate; Primary Ab: 1μg/ml Rabbit Anti-Rat TNFRSF1A Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



Immunohistochemistry

Image 3. Figure.DAB staining on IHC-P. Samples: Rat Tissue