
antibodies - online.com







anti-MSTN antibody (AA 267-376)





Overview

| Quantity: | 100 μL |
|----------------------|--|
| Target: | MSTN |
| Binding Specificity: | AA 267-376 |
| Reactivity: | Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This MSTN antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC) |

Product Details


| Immunogen: | MSTN (Arg267-Ser376) |
|---------------|--|
| Isotype: | IgG |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against MSTN. It has been selected for its ability to recognize MSTN in immunohistochemical staining and western blotting. |
| Purification: | Antigen-specific affinity chromatography |

Target Details

| Target: | MSTN |
|-------------|--|
| Abstract: | MSTN Products |
| Background: | Alternative Names: GDF8, Growth Differentiation Factor 8 |

Application Details

| 10 10 10 10 10 10 10 10 10 10 10 10 10 1 | |
|--|---|
| Application Notes: | Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. |
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated |
| | thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious |
| | degradation and precipitation were observed. The loss rate is less than 5% within the expiration |
| | date under appropriate storage condition. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | Lot specific |
| Buffer: | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. |
| Preservative: | Sodium azide |
| Precaution of Use: | WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or |
| | eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a |
| | physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute |
| | azide-containing compounds in running water before discarding to avoid accumulation of |
| | potentially explosive deposits in lead or copper plumbing. |
| Handling Advice: | Avoid repeated freeze-thaw cycles. |
| Storage: | 4 °C |
| Storage Comment: | Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months. |
| Expiry Date: | 12 months |
| | |




Western Blotting

Image 1.

Western Blotting

Image 2. Western Blot; Sample: Human Liver lysate; Primary Ab: 1μg/ml Rabbit Anti-Rat MSTN Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



22-

Immunohistochemistry

Image 3. Used in DAB staining on fromalin fixed paraffinembedded liver tissue

Please check the product details page for more images. Overall 4 images are available for ABIN1174892.