antibodies -online.com










Overview

Overview	
Quantity:	100 μL
Target:	LY9 (CD229)
Binding Specificity:	AA 2-212
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LY9 antibody is un-conjugated
Application:	Western Blotting (WB), Immunocytochemistry (ICC), Immunohistochemistry (IHC)
Product Details	
Immunogen:	LY9 (Val2-Thr212)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against LY9. It has been selected for its ability to recognize LY9 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography
Target Details	
Target:	LY9 (CD229)
Alternative Name:	Lymphocyte Antigen 9 (LY9) (CD229 Products)
Background:	Alternative Names: CD229, LY9, SLAMF3, Hly9, Mly9, SLAMF3, Cell surface molecule Ly-9,


Target Details

Signaling lymphocytic activation molecule 3
BCR Signaling
 Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.
The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
For Research Use only
Liquid
Lot specific
PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Sodium azide
WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Avoid repeated freeze-thaw cycles.
4 °C
Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.
12 months




Western Blotting

Image 1.



Western Blotting

Image 2. Western Blot; Sample: Human HepG2 cell lysate; Primary Ab: 3μg/ml Rabbit Anti-Human LY9 Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



Immunohistochemistry

Image 3. Figure.DAB staining on IHC-P. Samples: Human Tissue