

Datasheet for ABIN1175142  
**anti-CD3D antibody (AA 22-103)**



[Go to Product page](#)

**2** Images

## Overview

Quantity:	100 µL
Target:	CD3D
Binding Specificity:	AA 22-103
Reactivity:	Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CD3D antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP)

## Product Details

Purpose:	Polyclonal Antibody to Cluster Of Differentiation 3d (CD3d)
Immunogen:	RPB872Po01Recombinant Cluster Of Differentiation 3d (CD3d)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against CD3d. It has been selected for its ability to recognize CD3d in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

## Target Details

Target:	CD3D
---------	------

## Target Details


Alternative Name:	CD3d ( <a href="#">CD3D Products</a> )
Background:	CD3-DELTA, T3D, T-Dell Surface Glycoprotein CD3 Delta Chain, T-cell receptor T3 delta chain
Pathways:	<a href="#">TCR Signaling</a> , <a href="#">CXCR4-mediated Signaling Events</a>

## Application Details

Application Notes:	Western blotting: 0.5-2 µg/mL, Immunohistochemistry: 5-20 µg/mL, Immunocytochemistry: 5-20 µg/mL, Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Concentration:	500 µg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	12 months



Western Blotting

Image 1.



Immunohistochemistry

Image 2. DAB staining on IHC-P; Samples: Porcine Spleen Tissue