antibodies -online.com





anti-FURIN antibody (AA 105-221)

3 Images



Go to Product page

_							
0	V	е	r١	/1	е	V	I


Background:

Quantity:	100 μL
Target:	FURIN
Binding Specificity:	AA 105-221
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FURIN antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Immunogen:	FUR (Thr105-lle221)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against FUR. It has been selected for its ability to recognize FUR in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography
Target Details	
Target:	FURIN
Alternative Name:	Furin (FUR) (FURIN Products)

Alternative Names: PACE, PCSK3, SPC1, Paired Basic Amino Acid Cleaving Enzyme, Dibasic-


Target Details

rarget Details			
	processing enzyme		
Pathways:	Notch Signaling, Neurotrophin Signaling Pathway		
Application Details			
Application Notes:	 Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. 		
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	Lot specific		
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.		
Preservative:	Sodium azide		
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.		
Handling Advice:	Avoid repeated freeze-thaw cycles.		
Storage:	4 °C		
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.		
Expiry Date:	12 months		




Western Blotting

Image 1.



Immunohistochemistry

Image 2. DAB staining on IHC-P; Samples: Human Liver Tissue



Immunohistochemistry

Image 3. Used in DAB staining on fromalin fixed paraffinembedded Liver tissue