

# Datasheet for ABIN1176210 anti-HVEM antibody (AA 45-262)

# 2 Images



Go to Product page

$\sim$				
( )\	10	rVI	161	٨
$\cup$	ノロ	V	$\Box$	Λ

Quantity:	100 μL
Target:	HVEM (TNFRSF14)
Binding Specificity:	AA 45-262
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HVEM antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP),
	Immunocytochemistry (ICC)
Product Details	
Purpose:	Polyclonal Antibody to Tumor Necrosis Factor Receptor Superfamily, Member 14 (TNFRSF14)
Immunogen:	RPD926Mu01Recombinant Tumor Necrosis Factor Receptor Superfamily, Member 14
	(TNFRSF14)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against TNFRSF14. It has been selected for
	its ability to recognize TNFRSF14 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	
Target:	HVEM (TNFRSF14)


## Target Details

Alternative Name:	TNFRSF14 (TNFRSF14 Products)
Target Type:	Viral Protein
Background:	CD270, TR2, ATAR, HVEA, HVEM, LIGHTR, Herpesvirus Entry Mediator, Herpes virus entry
	mediator A, Tumor necrosis factor receptor-like 2
Pathways:	Production of Molecular Mediator of Immune Response, Cancer Immune Checkpoints
Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-
	20 μg/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	date under appropriate storage condition.  For Research Use only
Restrictions: Handling	
Handling	For Research Use only
Handling Format:	For Research Use only Liquid
Handling Format: Concentration:	For Research Use only  Liquid  500 μg/mL
Handling Format: Concentration: Buffer:	For Research Use only  Liquid  500 μg/mL  PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Handling  Format:  Concentration:  Buffer:  Preservative:	For Research Use only  Liquid  500 μg/mL  PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.  Sodium azide
Handling  Format:  Concentration:  Buffer:  Preservative:	For Research Use only  Liquid  500 μg/mL  PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.  Sodium azide  WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.
Handling  Format:  Concentration:  Buffer:  Preservative:	For Research Use only  Liquid  500 μg/mL  PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.  Sodium azide  WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.  Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or
Handling  Format:  Concentration:  Buffer:  Preservative:	For Research Use only  Liquid  500 µg/mL  PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.  Sodium azide  WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.  Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a
Handling  Format:  Concentration:  Buffer:  Preservative:	For Research Use only  Liquid  500 μg/mL  PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.  Sodium azide  WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.  Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute
Handling  Format:  Concentration:  Buffer:  Preservative:	Eiquid  500 μg/mL  PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.  Sodium azide  WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.  Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of
Handling  Format:  Concentration:  Buffer:  Preservative:  Precaution of Use:	Eiquid  500 µg/mL  PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.  Sodium azide  WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.  Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling  Format:  Concentration:  Buffer:  Preservative:  Precaution of Use:  Handling Advice:	For Research Use only  Liquid  500 μg/mL  PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.  Sodium azide  WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.  Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.  Avoid repeated freeze-thaw cycles.

Expiry Date:


12 months

#### **Images**



### **Western Blotting**

Image 1.



#### **Immunohistochemistry**

**Image 2.** Figure.DAB staining on IHC-P. Samples: Mouse Tissue