

Datasheet for ABIN1176764

anti-LCAT antibody (AA 290-433) (Biotin)





\sim		
	rview	
() (I V I C. V V	


Overview	
Quantity:	200 μL
Target:	LCAT
Binding Specificity:	AA 290-433
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LCAT antibody is conjugated to Biotin
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Purpose:	Biotin-Linked Polyclonal Antibody to Lecithin Cholesterol Acyltransferase (LCAT)
Immunogen:	Recombinant LCAT expressed in E.coli.
	The antibody is a rabbit polyclonal antibody raised against LCAT conjugated to biotin.
Sequence:	MGHHHHHHSG SEF-I STPSFNYTGR DFQRFFADLH FEEGWYMWLQ SRDLLAGLPA
	PGVEVYCLYG VGLPTPRTYI YDHGFPYTDP VGVLYEDGDD TVATRSTELC GLWQGRQPQP
	VHLLPLHGIQ HLNMVFSNLT LEHINAILLG AYRQGPPASP TAS
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against LCAT. It has been selected for its
	ability to recognize LCAT in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse, Rat

Antigen-specific affinity chromatography followed by Protein A affinity chromatography
LCAT
Lecithin Cholesterol Acyltransferase (LCAT Products)
Phosphatidylcholine-sterol O-acyltransferase, Phospholipid-cholesterol acyltransferase
Lipid Metabolism
Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100 Immunocytochemistry: 5-20 μg/mL,1:25-100 Optimal working dilutions must be determined by end user.
The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
For Research Use only
Liquid
500 μg/mL
PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Sodium azide
WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Avoid repeated freeze/thaw cycles
4 °C,-20 °C

Handling

Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	12 months
les s d s s	

Images



Western Blotting

Image 1.