# antibodies -online.com





# anti-ITGB1 antibody (APC)

2 Images

3

**Publications** 



Go to Product page

#### Overview

Quantity:	100 tests
Target:	ITGB1
Reactivity:	Human, Pig, Dog
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This ITGB1 antibody is conjugated to APC
Application:	Flow Cytometry (FACS)

### Product Details

1 Toddet Details	
Immunogen:	Raji Burkitt's lymphoma cell line
Clone:	MEM-101A
Isotype:	lgG1
Specificity:	The antibody MEM-101A reacts with an extracellular epitope of CD29 (integrin beta1 chain), a 130 kDa single chain type I glycoprotein expressed as a heterodimer (non-covalently associated with the integrin alpha subunits 1-6). CD29 is broadly expressed on majority of hematopoietic and non-hematopoietic cells (leukocytes, platelets, fibroblasts, endothelial cells, epithelial cells and mast cells).
No Cross-Reactivity:	Mouse
Cross-Reactivity (Details):	Human, Porcine, Canine (Dog)
Purification:	Purified antibody is conjugated with activated allophycocyanin (APC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion

chron	nator	ıran	hν	,
CHILOH	παιος	jιαρ	піу	


# Target Details

rarget Details	
Target:	ITGB1
Alternative Name:	CD29 (ITGB1 Products)
Background:	Integrin subunit beta 1,CD29 (beta1 integrin subunit, GPIIa) forms non-covalently linked heterodimers with at least 6 different alpha chains (alpha1-alpha6, CD49a-f) determining the binding properties of beta1 (VLA) integrins. These integrins mediate cell adhesion to collagen, fibronectin, laminin and other extracellular matrix (ECM) components. This interaction hinders cell death, whereas disruption of anchorage to ECM leads to apoptosis. Decreased expression of most beta1 integrins correlates with acquiring multidrug resistance of tumour cells during selection in presence of antitumour drug. In platelets, translocation of intracellular pool of beta1 integrins to the plasma membrane following thrombin stimulation. These integrins are also upregulated in leukocytes during emigration and extravascular migration and appear to be critically involved in regulating the immune cell trafficking from blood to tissue, as well as in regulating tissue damage and disease symptoms related to inflammatory bowel disease. Through a beta1 integrin-dependent mechanism, fibronectin and type I collagen enhance cytokine secretion of human airway smooth muscle in response to IL-1beta.,Integrin β,1 chain, ITGB1, VLAB, MDF2, FNRB, GPIIA, MSK12
Gene ID:	3688
UniProt:	P05556
Pathways:	Cell-Cell Junction Organization, Regulation of G-Protein Coupled Receptor Protein Signaling, CXCR4-mediated Signaling Events, Signaling of Hepatocyte Growth Factor Receptor, Integrin Complex, SARS-CoV-2 Protein Interactome
Application Details	
Application Notes:	Flow cytometry: The reagent is designed for analysis of human blood cells using 10 $\mu$ L reagent / 100 $\mu$ L of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.
Comment:	The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.

Application Details	
Restrictions:	For Research Use only
Handling	
Reconstitution:	No reconstitution is necessary.
Buffer:	Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Do not freeze.  Avoid prolonged exposure to light.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.
Publications	
Product cited in:	Miksiunas, Rucinskas, Janusauskas, Labeit, Bironaite: "Histone Deacetylase Inhibitor
	Suberoylanilide Hydroxamic Acid Improves Energetic Status and Cardiomyogenic
	Differentiation of Human Dilated Myocardium-Derived Primary Mesenchymal Cells." in:
	International journal of molecular sciences, Vol. 21, Issue 14, (2020) (PubMed).
	Plánka, Necas, Srnec, Rauser, Starý, Jancár, Amler, Filová, Hlucilová, Kren, Gál: "Use of allogenic stem cells for the prevention of bone bridge formation in miniature pigs." in: <b>Physiological</b>

research / Academia Scientiarum Bohemoslovaca, Vol. 58, Issue 6, pp. 885-93, (2010) ( PubMed).

Símová, Klíma, Cermak, Sourková, Andera: "Arf and Rho GAP adapter protein ARAP1 participates in the mobilization of TRAIL-R1/DR4 to the plasma membrane." in: Apoptosis: an international journal on programmed cell death, Vol. 13, Issue 3, pp. 423-36, (2008) (PubMed).



# **Flow Cytometry**

**Image 1.** Flow cytometry analysis (surface staining) of human peripheral blood with anti-human CD29 (MEM-101A) APC.

# **Western Blotting**

**Image 2.** Western Blotting analysis (non-reducing conditions) of isolated peripheral blood lymphocytes of various species using anti-CD29 (MEM-101A). Lane 1: lysate of human PBL Lane 2: lysate of canine PBL Lane 3: lysate of porcine PBL