

Datasheet for ABIN126717

anti-beta Amyloid antibody (C-Term, N-Term) (Biotin)





Go to Product page

_				
()	ve.	rv/	101	Λ

Quantity:	0.1 mg		
Target:	beta Amyloid (Abeta)		
Binding Specificity:	AA 1-40, C-Term, N-Term		
Reactivity:	Human		
Host:	Mouse		
Clonality:	Monoclonal		
Conjugate:	This beta Amyloid antibody is conjugated to Biotin		
Application:	Western Blotting (WB), Immunofluorescence (IF), Enzyme Immunoassay (EIA)		
Product Details			
Immunogen:	C-terminal peptide conjugated to KLH		
Clone:	5C3		
Isotype:	lgG1		
Specificity:	This antibody specifically interacts with the C-Terminus of beta-Amyloid (1-40) and does not		
	crossreact with beta-Amyloid (1-42).		
Purification:	Subsequent Thiophilic Adsorption and Size Exclusion Chromatography.		
Target Details			
Target:	beta Amyloid (Abeta)		
Alternative Name:	Amyloid beta (Abeta Products)		

Target Details

Expiry Date:

rarget Details			
Background:	The beta-amyloid peptide (beta A4), proteolytically released from the amyloid precursor protein		
	(APP), is the principal component of senile plaques in Alzheimer's disease. Cleavage of APP by		
	alpha-secretase or alternatively by beta-secretase leads to generation and extracellular release		
	of soluble APP peptides, S-APP-alpha and S-APP-beta, respectively, and the retention of		
	corresponding membrane-anchored C-terminal fragments, C83 and C99. Subsequent		
	processing of C83 by gamma-secretase yields P3 peptides. This is the major secretory pathway		
	and is nonamyloidogenic. Alternatively, presenilin/nicastrin-mediated gamma-secretase		
	processing of C99 releases the amyloid beta proteins, amyloid-beta 40 (Abeta40) and amyloid-		
	beta 42 (Abeta42), major components of amyloid plaques, and the cytotoxic C-terminal		
	fragments, gamma-CTF(50), gamma-CTF(57) and gamma-CTF(59). Synonyms: Amyloid beta		
	peptide		
Pathways:	Inflammasome		
Application Details			
Application Notes:	ELISA: 0.1 μg/mL. Immunocytochemistry: 1-10 μg/mL. Western blot: 1 μg/mL for HRPO/ECL		
	detection. Recommended blocking buffer: Casein/Tween 20 based blocking and blot		
	incubationbuffer.		
	Other applications not tested.		
	Optimal dilutions are dependent on conditions and should be determined by the user.		
Restrictions:	For Research Use only		
Handling			
Concentration:	0.5 mg/mL		
Buffer:	PBS containing 0.09 % Sodium Azide as preservative.		
Preservative:	Sodium azide		
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which		
	should be handled by trained staff only.		
Storage:	4 °C/-80 °C		
Storage Comment:	Store the antibody (aliquote in liquid nitrogen) at -80 °C. Avoid repeated freezing and thawing.		
	Thaw aliquots at 37 °C. Thawed aliquots may be stored at 2-8 °C up to 3 months.		
	Shelf life: one year from despatch.		

12 months



Western Blotting

Image 1.