# antibodies -online.com







# anti-RNF2 antibody (AA 189-201)





$\sim$					
	1//	r۱.	/ I	$\triangle$	٨

Quantity:	100 μg
Target:	RNF2
Binding Specificity:	AA 189-201
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This RNF2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA
Product Details	

	with a synthetic peptide corresponding aa 189-201 of human RING1B protein.
Isotype:	lgG
Characteristics:	Concentration Definition: by UV absorbance at 280 nm

## **Target Details**

Target:	RNF2
Alternative Name:	RING1B (RNF2 Products)
Background:	RING1B (also known as BAP1, DING, Polycomb-M33 interacting protein Ring1B, Ring finger protein 1b, Ring finger protein 2 and RNF2) is one of the PcG proteins. The polycomb group
	(PcG) of proteins form the multiprotein complexes that are important for the transcription

repression of various genes involved in development and cell proliferation. It has been shown to interact with, and suppress the activity of, transcription factor CP2 (TFCP2/CP2). Studies of the mouse counterpart suggested the involvement of this gene in the specification of anterior-posterior axis, as well as in cell proliferation in early development. This protein was also found to interact with huntingtin interacting protein 2 (HIP2), a ubiquitin-conjugating enzyme that possesses ubiquitin ligase activity.

Synonyms: BAP1 antibody, DING antibody, DinG protein antibody, RING finger protein BAP1 antibody, E3 ubiquitin protein ligase RING 2 antibody, Huntingtin interacting protein 2 interacting protein 3 antibody, RNF 2 antibody

Gene ID: 6045, 6005747

UniProt: Q99496

### **Application Details**

Application Notes: Anti-RING1B purified antibody has been tested for use in ELISA and by western blot. Specific


conditions for reactivity should be optimized by the end user. Expect a band approximately 38 kDa in size corresponding to RING1B by western blotting in the appropriate cell lysate or

extract.

Restrictions: For Research Use only

#### Handling

Format:	Liquid
Concentration:	1.0 mg/mL
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C







DAPI Nuclear staining

Tubulin antibody staining (cytoplasm)





Rnf2 antibody

Merged picture (DAPI: blue; Tubulin: red Rnf2: green)

#### **Western Blotting**

Image 1. Western blot using Affinity Purified anti-RING1B antibody shows detection of a 38 kDa band corresponding to human RING1B in 3T3 (lane 1), U937 (lane 2), Jurkat (lane 3), mouse brain (lane 4) and CHO-K1 (lane 5) cell lysates. Approximately 20 µg of lysate was run on a SDS-PAGE and transferred onto nitrocellulose followed by reaction with a 1:500 dilution of anti-RING1B antibody incubated at room temper-ature. Signal was detected using standard techniques.

#### **Immunofluorescence**

Image 2. Immunofluorescence Microscopy of Goat anti-RING1B antibody. Tissue: human HeLa cells. Fixation: methanol and blocked with 0.2% fish scale gelatin for 1 hour at 25°C. Antigen retrieval: not required. Primary antibody: RING1B antibody at 1:300 for 20 minutes at 25°C. Secondary antibody: Alexa488-conjugated Donkey anti-goat IgG secondary antibody at 1:500 for 45 min at RT. Localization: RING1B is nuclear and occasionally cytoplasmic. Staining: RING1B (RNF2) as green signal, Tubulin cytoplasm staining red, and DAPI (blue) nuclear counterstain.