.-online.com antibodies

Datasheet for ABIN1302413 anti-PDGFRB antibody

1 Image

1 Publication



Overview

Quantity:	0.1 mg
Target:	PDGFRB
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This PDGFRB antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunocytochemistry (ICC)

Product Details

Immunogen:	CD140b-transfected NIH 3T3 cells
Clone:	18A2
Isotype:	lgG1
Specificity:	The monoclonal antibody 18A2 recognizes an extracellular epitope of CD140b / PDGF-RB, the 180-190 kDa beta chain of platelet-derived growth factor receptor, which is widely expressed on a variety of mesenchymal-derived cells and plays pro-proliferative roles.
Cross-Reactivity (Details):	Human
Purification:	Purified by protein-A affinity chromatography.
Purity:	> 95 % (by SDS-PAGE)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN1302413 | 01/29/2024 | Copyright antibodies-online. All rights reserved.

Target Details

Target:	PDGFRB
Abstract:	PDGFRB Products
Background:	Platelet derived growth factor receptor beta,CD140b / PDGF-RB (platelet-derived growth factor receptor beta) is a cell surface receptor for members of platelet-derived growth factor family, whose intracellular part contains a tyrosine kinase domain. CD140b forms homodimers, or heterodimerizes with CD140a / PDGF-RA. Whereas CD140a can have both pro-proliferative or anti-proliferative effects, the CD140b induces in various cell types their proliferation and migration. CD140b has also developmental roles in the cardiovascular system and is preferentially expressed on some tumours such as medulloblastoma.,PDGFRB, PDGFR1
Gene ID:	5159
UniProt:	Q59F04
Pathways:	Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Inositol Metabolic Process, Glycosaminoglycan Metabolic Process, Smooth Muscle Cell Migration, Platelet-derived growth Factor Receptor Signaling


Application Details

Application Notes:	Flow cytometry: Recommended dilution: 1-4 µg/mL.
Restrictions:	For Research Use only
Handling	
Concentration:	1 mg/mL
Buffer:	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Do not freeze.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Do not freeze.

Product cited in:

Fekete, Rojewski, Fürst, Kreja, Ignatius, Dausend, Schrezenmeier: "GMP-compliant isolation and large-scale expansion of bone marrow-derived MSC." in: **PLoS ONE**, Vol. 7, Issue 8, pp. e43255, (2012) (PubMed).

Images



Flow Cytometry

Image 1. Flow cytometry (surface staining) of CD140b / PDGF-RB in human CD140b / PDGF-RB stable transfectants with anti-CD140b (18A2) purified GAM-APC.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN1302413 | 01/29/2024 | Copyright antibodies-online. All rights reserved.