


# Datasheet for ABIN1302424

# anti-FOXP3 antibody





#### Go to Product page

$\sim$			
( )	1/0	r\/I	$\Theta M$
$\cup$	$\vee \subset$	I V I	lew

Quantity:	100 μg	
Target:	FOXP3	
Reactivity:	Human, Mouse	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This FOXP3 antibody is un-conjugated	
Application:	Western Blotting (WB), Flow Cytometry (FACS)	

#### **Product Details**

Purpose:	Anti-FoxP3 Purified
Immunogen:	Full-length His-tagged recombinant murine FoxP3
Clone:	3G3
Isotype:	IgG1 kappa
Specificity:	The mouse monoclonal antibody 3G3 recognizes N-terminal region of FoxP3, a 47-55 kDa transcription factor (intracellular antigen), which is the master regulator in the development and function of regulatory T cells.
Cross-Reactivity (Details):	Human, Mouse
Purification:	Purified by protein-A affinity chromatography.
Purity:	> 95 % (by SDS-PAGE)

# **Target Details**


Target:	FOXP3	
Alternative Name:	FoxP3 (FOXP3 Products)	
Background:	Forkhead box P3,FoxP3 (forkhead box protein 3), a highly conserved forkhead/winged-helix	
	transcription factor, plays a crucial role in maintaining immune homeostasis by governing the	
	development and function of regulatory T cells. It is constitutively expressed at high level in	
	CD25+ CD4+ Treg cells and at low level in a CD25- CD4+ Treg cell subset. Defects in gene	
	encoding FoxP3 protein cause the scurfy phenotype in mice, and in human the IPEX syndrome	
	(immune dysfunction, polyendocrinopathy, enteropathy, X-linked syndrome), also known as X-	
	linked autoimmunity-allergic dysregulation (XLAAD) syndrome.,Scurfin, JM2, AIID, IPEX, PIDX,	
	XPID, DIETER	
Gene ID:	50943	
UniProt:	Q9BZS1	
Pathways:	Chromatin Binding, Regulation of Leukocyte Mediated Immunity, Positive Regulation of	
	Immune Effector Process, Production of Molecular Mediator of Immune Response, Activated 7	
	Cell Proliferation	
A 11 D		
Application Details		
Application Notes:	Western blotting: Recommended dilution: 2 μg/mL.	
	Flow cytometry: Recommended dilution: 1-4 $\mu$ g/mL. Intracellular staining.	
Restrictions:	For Research Use only	
Handling		
Concentration:	1 mg/mL	
Buffer:	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
	should be handled by trained staff only.	
Handling Advice:	Do not freeze.	
Storage:	4 °C	
Storage Comment:	Store at 2-8°C. Do not freeze.	

Product cited in:

Law, Hirschkorn, Owen, Biswas, Norris, Lanteri: "The importance of Foxp3 antibody and fixation/permeabilization buffer combinations in identifying CD4+CD25+Foxp3+ regulatory T cells." in: **Cytometry. Part A: the journal of the International Society for Analytical Cytology**, Vol. 75, Issue 12, pp. 1040-50, (2009) (PubMed).

Gavin, Torgerson, Houston, DeRoos, Ho, Stray-Pedersen, Ocheltree, Greenberg, Ochs, Rudensky: "Single-cell analysis of normal and FOXP3-mutant human T cells: FOXP3 expression without regulatory T cell development." in: **Proceedings of the National Academy of Sciences of the United States of America**, Vol. 103, Issue 17, pp. 6659-64, (2006) (PubMed).

### **Images**



#### Immunofluorescence

**Image 1.** Intracellular staining of human peripheral blood cells (gated on CD4+ cells) with anti-FoxP3 (clone 3G3).