antibodies -online.com





anti-EVI2B antibody

3 Images



Go to Product page

Overview


Quantity:	0.1 mg
Target:	EVI2B
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This EVI2B antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunoprecipitation (IP)

Product Details

Immunogen:	Raji cells
Clone:	MEM-216
Isotype:	IgG1 kappa
Specificity:	The mouse monoclonal antibody MEM-216 recognizes an extracellular epitope of CD361 / EVI2B, almost uncharacterized type I transmembrane protein with broad leukocyte expression, mostly in myeloid and B cells.
Cross-Reactivity (Details):	Human
Purification:	Purified by protein-A affinity chromatography.
Purity:	> 95 % (by SDS-PAGE)


Target Details

EVI2B
CD361 (EVI2B Products)
Ecotropic viral integration site 2B,CD361, also known as EVI2B (ecotropic viral integration site
2B) or EVDB, is a poorly characterized type I transmembrane protein, expressed from one of
three genes embedded in intron 27b of the neurofibromatosis type 1 (NF1) gene. The DNA
strand that is transcribed to produce CD361 is the complementary one to the strand encoding
NF1. Murine homolog to human CD361 is associated with ecotropic viral insertions, which have
been implicated in the expression of murine myeloid leukemias. CD361 has been also reported
to be involved in melanocyte and keratinocyte differentiation. However, it is expressed mainly i
peripheral blood and bone marrow.,EVI2B, EVDB
2124
P34910
Flow cytometry: Recommended dilution: 1-4 µg/mL, Positive control, Positive control: Raji,
Daudi, HL-60 cells, peripheral blood lymphcocytes (strongly positive on CD19+ cells), negative
control: Jurkat, U-937 cells.
For Research Use only
1 mg/mL
Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Sodium azide
This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
should be handled by trained staff only.
Do not freeze.
4 °C
Store at 2-8°C. Do not freeze.




Flow Cytometry

Image 1. Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD361 (MEM-216) purified antibody (concentration in sample 4 μ g/mL, GAM APC).



Flow Cytometry

Image 2. Flow cytometry analysis of CD19+ peripheral blood leukocytes using anti-CD361 antibody (MEM-216).



Flow Cytometry

Image 3. Separation of human neutrophil granulocytes (red-filled) from lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD361 (MEM-216) purified antibody (concentration in sample 4 μ g/mL, GAM APC).