

Datasheet for ABIN1302471

**anti-CD161 antibody****2** Images**4** Publications[Go to Product page](#)

## Overview

Quantity:	0.1 mg
Target:	CD161 (KLRB1)
Reactivity:	Rat
Host:	Mouse
Clonality:	Monoclonal
Application:	Flow Cytometry (FACS), Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (Frozen Sections) (IHC (fro)), Radioimmunoassay (RIA)

## Product Details

Immunogen:	Splenic cells purified from the LEW rat
Clone:	10-78
Isotype:	IgG1 kappa
Specificity:	The mouse monoclonal antibody 10/78 recognizes CD161, an approximately 30 kDa type II transmembrane C-type lectin receptor, expressed on the plasma membrane of NK cells, dendritic cells, activated monocytes and a subset of T cells as a disulphide-linked homodimer. A common extracellular epitope on rat CD161a and b isoforms is detected.
Cross-Reactivity (Details):	Rat
Purification:	Purified by protein-A affinity chromatography.
Purity:	> 95 % (by SDS-PAGE)

## Target Details

Target:	CD161 (KLRB1)
Alternative Name:	CD161 ( <a href="#">KLRB1 Products</a> )
Background:	<p>Killer cell lectin-like receptor subfamily B, memb,CD161, also known as Nkrp1 (natural killer receptor protein 1) or Klrb1 (killer cell lectin-like receptor subfamily b member 1), is a disulphide-linked homodimeric receptor, which is involved in regulation of NK cell and NKT cell function. It is expressed on rat NK cells, subset of T cells, dendritic cells, and activated monocytes.</p> <p>Although human CD161 is expressed as one isoform, the rat CD161 has three isoforms, referred to as CD161a, b, and c. These proteins contain C-terminal C-type lectin extracellular domain, a transmembrane domain, and N-terminal intracellular domain, which contains ITIM motif, such as CD161b, and displays inhibitory function, or does not contain ITIM motif, thus also not the inhibitory function, such as CD161a.,NKRP1A, Klrb</p>
Gene ID:	689817
UniProt:	<a href="#">Q0ZUP0</a>

## Application Details

Application Notes:	Flow cytometry: Recommended dilution: 1-4 µg/mL.
Restrictions:	For Research Use only

## Handling

Concentration:	1 mg/mL
Buffer:	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	<b>Do not freeze.</b>
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Do not freeze.

## Publications


Product cited in:	Stephens, Barclay, Mason: "Phenotypic characterization of regulatory CD4+CD25+ T cells in rats." in: <b>International immunology</b> , Vol. 16, Issue 2, pp. 365-75, (2004) ( <a href="#">PubMed</a> ).
-------------------	---

May, Dorris, Satumtira, Iqbal, Rehman, Lightfoot, Taurog: "CD8 alpha beta T cells are not essential to the pathogenesis of arthritis or colitis in HLA-B27 transgenic rats." in: **Journal of immunology (Baltimore, Md. : 1950)**, Vol. 170, Issue 2, pp. 1099-105, (2003) ([PubMed](#)).

Tliba, Chauvin, Le Vern, Boulard, Sbille: "Evaluation of the hepatic NK cell response during the early phase of Fasciola hepatica infection in rats." in: **Veterinary research**, Vol. 33, Issue 3, pp. 327-32, (2002) ([PubMed](#)).

Kraus, Lambracht, Wonigeit, Hünig: "Negative regulation of rat natural killer cell activity by major histocompatibility complex class I recognition." in: **European journal of immunology**, Vol. 26, Issue 11, pp. 2582-6, (1997) ([PubMed](#)).

Images



Flow Cytometry

**Image 1.** Flow cytometry surface staining pattern of rat splenocyte suspension stained using anti-rat CD161 (10/78) purified antibody (concentration in sample 0,5 µg/mL) GAM APC.

Flow Cytometry

**Image 2.** Separation of rat CD161 positive splenocytes (red-filled) from CD161 negative splenocytes (black-dashed) in flow cytometry analysis (surface staining) of rat splenocyte suspension stained using anti-rat CD161 (10/78) purified antibody (concentration in sample 0,5 µg/mL) GAM APC.